

Acute pancreatitis: recent advances through randomised trials

Sven M van Dijk,¹ Nora D L Hallensleben,² Hjalmar C van Santvoort,³ Paul Fockens,⁴ Harry van Goor,⁵ Marco J Bruno,² Marc G Besselink,¹ for the Dutch Pancreatitis Study Group

¹Department of Surgery, Amsterdam Gastroenterology and Metabolism, Academic Medical Center, Amsterdam, The Netherlands

²Department of Gastroenterology & Hepatology, Erasmus University Medical Center, Rotterdam, The Netherlands

³Department of Surgery, St. Antonius Hospital, Nieuwegein, The Netherlands

⁴Department of Gastroenterology & Hepatology, Amsterdam Gastroenterology and Metabolism, Academic Medical Center, Amsterdam, The Netherlands

⁵Department of Surgery, Radboud University Medical Center, Nijmegen, The Netherlands

Correspondence to

Marc G Besselink, Department of Surgery, Amsterdam Gastroenterology and Metabolism, AMC Amsterdam, 22660, Amsterdam, The Netherlands; m.g.besselink@amc.uva.nl

SMD and NDH contributed equally.

Received 20 April 2017

Revised 5 July 2017

Accepted 7 July 2017

Published Online First

24 August 2017

ABSTRACT

Acute pancreatitis is one of the most common GI conditions requiring acute hospitalisation and has a rising incidence. In recent years, important insights on the management of acute pancreatitis have been obtained through numerous randomised controlled trials. Based on this evidence, the treatment of acute pancreatitis has gradually developed towards a tailored, multidisciplinary effort, with distinctive roles for gastroenterologists, radiologists and surgeons. This review summarises how to diagnose, classify and manage patients with acute pancreatitis, emphasising the evidence obtained through randomised controlled trials.

BACKGROUND

With over 26 000 hospital admissions in the UK each year, acute pancreatitis is among the most common GI conditions requiring acute hospitalisation.¹ The worldwide incidence of acute pancreatitis is rising, thus further increasing its burden on healthcare services.² Acute pancreatitis is an inflammatory process which causes a local and systemic inflammatory response syndrome (SIRS). Although the majority of patients have a mild disease course, around 20% will develop moderate or severe pancreatitis, with necrosis of the (peri)pancreatic tissue and/or (multiple-)organ failure (figure 1).³

Over the last decades, the treatment of acute pancreatitis has gradually developed towards a tailored, multidisciplinary approach, with a distinctive role for endoscopic, radiological and surgical treatment strategies. Much of the new evidence on the treatment of acute pancreatitis arises from randomised controlled trials (RCTs), which are universally considered as the reference standard for comparing treatment strategies in medicine. Random allocation of patients, if possible blinded, keeps all known and unknown variables constant except for the allocated treatment, thereby measuring the 'true' effect of the investigated treatment. In the past decade, numerous RCTs have had a great impact on the treatment of acute pancreatitis. This review provides an overview of current clinical practice concerning the diagnosis, classification and treatment of acute pancreatitis, while focussing on the outcomes of these RCTs.

METHODS

For this review, the most recent international evidenced-based guidelines on acute pancreatitis, the 2012 International Association of

Significance of this study

Early phase

- ▶ Primary management of acute pancreatitis consists of (early goal-directed) fluid resuscitation with Ringer's lactate and adequate pain control.
- ▶ Endoscopic retrograde cholangiography/endoscopic sphincterotomy should be performed urgently in case of concomitant cholangitis, should not be performed in predicted mild biliary pancreatitis and is controversial in predicted severe biliary pancreatitis.
- ▶ Prophylactic use of antibiotics or probiotics is not indicated.
- ▶ In patients with acute pancreatitis, regardless of severity, a normal oral diet can be started once the acute pain is resolving.
- ▶ Nasoenteric tube feeding is indicated only when sufficient oral intake is not reached after 3–5 days.

Beyond the early phase

- ▶ Cholecystectomy should be performed during the index admission for mild biliary pancreatitis.
- ▶ Intervention for infected necrotising pancreatitis should preferably be delayed until the phase of walled-off necrosis.
- ▶ The step-up approach, either endoscopic or surgical, is the preferred treatment of infected necrotising pancreatitis.

Aftercare

- ▶ In (presumed) idiopathic pancreatitis, a repeat abdominal ultrasound and ultimately endoscopic ultrasound may detect microlithiasis/sludge in up to half of patients, warranting cholecystectomy.
- ▶ Alcohol abstinence support programmes can prevent recurrent alcoholic pancreatitis.
- ▶ Attention should be paid to potential endocrine and exocrine insufficiency after necrotising pancreatitis.

Pancreatology/American Pancreatic Association (IAP/APA) guidelines,⁴ were used as the starting point. PubMed was searched for studies, specifically RCTs, published after the IAP/APA guideline using the following terms: Pancreatitis (MeSH Terms) OR



CrossMark

To cite: van Dijk SM, Hallensleben ND, van Santvoort HC, *et al.* *Gut* 2017;**66**:2024–2032.




Figure 1 Mortality rates of acute pancreatitis.

(acute pancreatitis (Title)). All articles regarding chronic pancreatitis and malignant disease were excluded. Two authors (SvD and NH) assessed all English articles concerning RCTs on adults published between June 2012 and February 2017. In total, 490 articles were found, of which all potential ‘practice changing’ RCTs were incorporated within this review. Additionally, relevant articles from the reference list of the included articles were reviewed as well as new, evidence-based guidelines on acute pancreatitis published after the IAP/APA guideline.

EARLY PHASE OF ACUTE PANCREATITIS

Diagnosis

According to the 2012 Revised Atlanta Classification, the diagnosis acute pancreatitis requires at least two of the three following criteria: (1) abdominal pain consistent with pancreatitis, (2) serum amylase and/or lipase of at least three times the upper limit of the normal value or (3) findings consistent with acute pancreatitis on imaging (contrast-enhanced CT (CECT), MRI or ultrasound).⁵ In case of typical clinical and laboratory findings, an additional CECT or other cross-sectional imaging is not required to confirm the diagnosis. The severity of acute pancreatitis is classified as mild, moderate or severe. Mild when there are no local or systemic complications present; moderate in case of local (eg, peripancreatic fluid collections) or systemic complications (eg, exacerbation of chronic disease) or transient organ failure (<48 hours) and severe in case of persistent organ failure (>48 hours).⁵

Aetiology

In Western countries, gallstones and/or biliary sludge are the most prevalent (approximately 40%–50%) cause of acute pancreatitis.^{6,7} With approximately 20% of cases, alcohol is the second most frequent cause of acute pancreatitis in most countries.^{6–8} Less frequent causes of acute pancreatitis include

medication, endoscopic retrograde cholangiopancreatography, hypercalcaemia, hypertriglyceridemia, surgery and trauma.

Determining the aetiology of acute pancreatitis is of importance, as it partly drives early management as well as the follow-up strategy. Standard work-up of acute pancreatitis includes medical history, physical examination, laboratory tests (liver enzymes, triglycerides, calcium) and transabdominal ultrasound. In 10%–25% of cases, the aetiology of the pancreatitis remains unclear. Idiopathic pancreatitis requires additional diagnostic work-up in the form of a repeat transabdominal ultrasound and ultimately an endoscopic ultrasound (EUS).⁴ Meta-analyses show that in around 61% of cases, an aetiology can be established by EUS. This includes the detection of micro-lithiasis or biliary sludge (41%), for which cholecystectomy is




Figure 2 Linear endoscopic ultrasound image of sludge in the gallbladder.

Recent advances in clinical practice

required to prevent recurrences (figure 2), but also other causes such as chronic pancreatitis or pancreatic tumours.⁹

Prediction of severity

Prediction of severity of acute pancreatitis is used to identify patients at low or high risk of developing complications. This is useful both for triaging patients for the proper level of monitoring and for stratification of patients for inclusion in RCTs. Multiple scoring systems are available that combine clinical and laboratory findings to determine the likelihood of a severe disease course: the Acute Physiology and Chronic Health Evaluation II (APACHE-II), Ranson score, modified Glasgow/Imrie score, SIRS criteria, Bedside Index for the Severity in Acute Pancreatitis and Harmless Acute Pancreatitis Score, while single laboratory parameters such as C reactive protein (CRP) can also be used.

Approximately 50% of all patients have predicted severe pancreatitis. Although only half of these will ultimately develop (moderately) severe pancreatitis, virtually none of the patients with predicted mild pancreatitis will do so. Mortality of predicted severe pancreatitis is approximately 10%, as compared with <1% in patients with predicted mild pancreatitis. The scoring systems are therefore primarily used to exclude the possibility of developing severe pancreatitis. Since the accuracy of the various predictive scoring systems is comparable,¹⁰ the IAP/APA guideline advises to use persistent (>48 hours) SIRS because of its relative simplicity.⁴

Treatment in the acute phase

As no curative therapy is currently available for acute pancreatitis, early treatment consists of supportive care which includes adequate fluid resuscitation and pain management.

Fluid resuscitation

Inflammation of the pancreas and the accompanying systemic inflammatory response leads to extravasation of fluid to the third space. In severe cases this may cause hypovolaemia, hypoperfusion and ultimately organ failure. To counteract this cascade, adequate fluid resuscitation is needed. Few RCTs studied the type of fluids. In critically ill patients in general, colloids are discouraged since there is no evidence to support their effectiveness whereas hydroxyethyl starch might even increase mortality.¹¹ The IAP/APA guideline therefore proposes crystalloids in the form of Ringer's lactate. This is based on one multi-centre RCT in 40 patients with acute pancreatitis that showed beneficial effects on CRP levels and SIRS with Ringer's lactate as compared with normal saline.^{4 12} This advantage of Ringer's lactate or other balanced fluids (eg, Plasma-Lyte) over normal saline in pancreatitis patients has however not been confirmed by larger RCTs. Further studies are needed as multiple RCTs in the general intensive care setting have so far failed to find better outcomes when using balanced fluids.^{13–16}

Five RCTs using different fluid resuscitation protocols have been conducted.^{12 17–20} Two of these RCTs in 76 and 115 patients with severe acute pancreatitis show that rapid, uncontrolled fluid resuscitation (10–15 mL/kg/h or until a haematocrit <35% within 48 hours) significantly worsened the rates of infections, abdominal compartment syndrome, the need for mechanical ventilation and even mortality.^{17 18}

One RCT in 60 patients with predicted mild pancreatitis demonstrated that aggressive fluid hydration with Ringer's lactate (20 mL/kg bolus followed by 3 mL/kg/h) compared with standard hydration with Ringer's lactate (10 mL/kg bolus

followed by 1.5 mL/kg/h) improved a composite end point (ie, 'clinical improvement within 36 hours').¹⁹ So, both too little as well as too much fluid administration in the acute phase of pancreatitis can be harmful, possibly dependent on the severity of the pancreatitis. As advised by the IAP/APA guideline, early goal-directed therapy with adequate monitoring might therefore remain the preferred solution.⁴

Early goal-directed therapy was the subject of one multi-centre RCT in 40 acute pancreatitis patients and one RCT in 200 patients with severe acute pancreatitis.^{12 20} The first RCT could not confirm superiority of early goal-directed therapy, but the incidence of SIRS in the entire RCT was very low, suggesting less severe pancreatitis patients. The second RCT showed reductions in the duration of mechanical ventilation and in the rates of multiple organ failure and mortality in the early goal-directed therapy group, especially if fresh frozen plasma was added as a resuscitation fluid. However, as baseline APACHE-II scores were significantly worse in the control group, suggesting non-balanced randomisation, further RCTs are needed.

It is difficult to draw conclusions based on these RCTs as multiple of the three main parameters of interest (fluid type, fluid protocol and resuscitation goals) differ between trials. Since no single parameter adequately reflects hydration status, it is therefore advised to observe trends in multiple parameters. On the nursing ward, target measures include a heart rate <120/min, a mean arterial pressure between 64 and 85 mm Hg and a urinary output of at least 0.5 mL/kg/h. Relevant laboratory findings include blood urea nitrogen, creatinine levels and especially haematocrit, which should stay between 35% and 44%.⁴

Pain management

Pain is the predominant symptom of acute pancreatitis and should be treated promptly and adequately. Frequent reassessment of pain scores and, if indicated, adjustment of analgesic types and/or dosages is needed to assure proper pain management. Several RCTs compared different types of analgesia in acute pancreatitis.^{21–28} A systematic review on opioid use in acute pancreatitis and a recent meta-analysis reported that the quality of the majority of these RCTs is low and no particular analgesic strategy is superior.^{29 30} As current evidence is limited, pain can be managed according to general state-of-the-art pain protocols.

Antibiotics and probiotics as prophylaxis

One of the most lethal complications of acute pancreatitis is secondary infection of pancreatic or peripancreatic necrosis.³¹ This is thought to occur as a result of bacterial translocation from the gut.³² Several double-blind RCTs failed to show a reduction of infection of (peri)pancreatic necrosis through the prophylactic use of antibiotics,^{33–35} as confirmed by meta-analyses.^{36 37} Antibiotics are therefore only indicated when infection is either proven or clinically suspected. In order to prevent bacterial translocation, attempts were made to influence the intestinal microbiome using probiotic bacteria. Two RCTs compared probiotic bacteria with placebo in 62 and 45 patients with severe acute pancreatitis and reported promising results.^{38 39} A subsequent multicentre RCT* in 296 patients with predicted severe pancreatitis, however, showed increased rates of mortality and non-occlusive mesenteric ischaemia in patients receiving probiotics.⁴⁰ Therefore, administration of probiotics is currently considered contraindicated in the treatment of (predicted) severe pancreatitis.

Nutrition

Enteral nutrition does not only provide adequate caloric intake, it may also improve clinical outcomes. It has been hypothesised that the combination of disturbed intestinal motility, bacterial overgrowth and increased permeability of the gut can lead to bacterial translocation, thus causing infection of pancreatic necrosis.^{32 41–45} Enteral nutrition may reduce translocation by stimulating intestinal motility, reducing bacterial overgrowth and thereby maintaining mucosal gut integrity.^{46 47} A Cochrane review involving eight RCTs confirmed this, showing reduced rates of infection, organ failure and mortality in 348 patients with acute pancreatitis receiving routine enteral nutrition as compared with routine total parenteral nutrition.⁴⁸

Furthermore, the timing of initiation of enteral nutrition could also be relevant. Some retrospective studies suggested that an early start of nasoenteric feeding significantly reduced infection rates.^{49–51} A multicentre RCT* in 208 patients with predicted severe pancreatitis, comparing very early nasojejunal feeding (<24 hours) with introduction of an oral diet after 72 hours (with on-demand nasojejunal feeding) showed no beneficial effects on infection rates or mortality. Importantly, in the control group, 69% of patients did not require a nasoenteral tube, thus avoiding potential patient discomfort.⁵² A second recent RCT, comparing early nasojejunal feeding (<24 hour) with no nutritional support in 214 patients, also failed to show benefits from early nutritional support.⁵³ Based on these RCTs, tube feeding in predicted severe pancreatitis can be limited to those patients who have insufficient oral caloric intake after 3–5 days.

It was previously believed that nasogastric feeding in acute pancreatitis would increase the risk of aspiration, and increase inflammation and pain as a result of stimulation of the pancreatic excretion. However, three RCTs found nasogastric feeding non-inferior to nasojejunal feeding.^{54–56} Consequently, both routes of enteral feeding are now considered feasible and safe.⁵⁷ In patients with (predicted) mild pancreatitis, three RCTs have shown that a normal oral diet can be resumed once the pain is decreasing.^{58–60}

Endoscopic retrograde cholangiography in biliary pancreatitis

In patients with acute biliary pancreatitis, (transient) obstruction at the level of Vater's ampulla is thought to initiate pancreatic inflammation. Persisting biliary obstruction may aggravate the disease course. Early biliary decompression and stone removal using endoscopic retrograde cholangiography (ERC) with endoscopic sphincterotomy (ES) has therefore been extensively studied as a potential intervention to improve clinical outcomes in biliary pancreatitis. Common bile duct (CBD) stones may, however, pass into the duodenum spontaneously, in which case ERC with ES might be redundant and even harmful. Several RCTs have shown that early ERC is not effective in patients with predicted mild pancreatitis,^{61–63} as the potential benefits do not outweigh the procedural risks.

Emergency ERC with ES is indicated within 24 hours after diagnosing acute biliary pancreatitis with concomitant cholangitis.^{4 64} Importantly, diagnosing cholangitis in patients also suffering from SIRS, as is commonly seen in the early phase of (predicted severe) biliary pancreatitis, can be challenging. The definition of cholangitis differs between trials and includes Charcot's triad,⁶¹ expert opinion⁶⁵ and the updated Tokyo guidelines (TG13) for acute cholangitis.⁶⁶ These definitions do not take into account the underlying cause, namely acute pancreatitis, and have low thresholds regarding the presence of inflammation and cholestasis. This may lead to overdiagnosing of acute

cholangitis in acute pancreatitis, potentially exposing patients to unnecessary ERC procedures.⁶⁶ Further research is needed to establish appropriate diagnostic criteria for cholangitis in acute biliary pancreatitis patients.

The use of routine (early) ERC with ES in predicted severe biliary disease course is controversial. Several RCTs,^{61–63 65 67 68} subsequent meta-analyses and guidelines provide conflicting advice on this issue.⁶⁹ A possible explanation for these discrepancies are the differences in the definitions used for biliary pancreatitis and cholangitis, as well as differences in patient populations and in timing and quality of the ERC. For example, in some studies, ERC was performed in non-biliary pancreatitis; in other studies, patients with cholangitis were randomised while these patients should always undergo ERC and sphincterotomy was not performed routinely. Recently, a multicentre RCT* comparing routine early ERC plus ES with conservative treatment in 232 patients with predicted severe acute biliary pancreatitis but without cholangitis was completed and results are awaited (the APEC-trial, ISRCTN97372133).⁷⁰

The use of EUS for the detection of CBD stones is emerging in biliary pancreatitis. The sensitivity and specificity of EUS for CBD stone detection are superior to both transabdominal ultrasound and serum markers. As ERC with ES is presumably most effective in patients with persisting CBD stones, an EUS-first strategy to establish the indication for ERC with ES in acute biliary pancreatitis patients could improve outcomes. A meta-analysis comparing an EUS-first strategy with ERC including one RCT in 140 patients showed promising results. Around 71% of ERC procedures could be avoided without a negative effect on the clinical course of the pancreatitis.^{71 72} A recent prospective cohort study supports these observations.⁷³ Furthermore, a decision tree analysis on cost-effectiveness demonstrated that, especially in patients with severe acute biliary pancreatitis, an EUS-first method is less costly than ERC only.⁷⁴ Further research is needed to confirm the effectiveness and feasibility of this relatively new strategy.

BEYOND THE EARLY PHASE

Imaging

The 1992 Atlanta Classification was a global consensus on how to define acute pancreatitis and how to classify the severity and local pancreatic complications.³¹ In 2012, the Atlanta classification was revised to better define the morphology of acute pancreatitis and (peri)pancreatic collections as seen on CECT.⁵ The revised Atlanta classification distinguishes interstitial oedematous pancreatitis from necrotising pancreatitis, wherein the latter is subdivided in parenchymal and peripancreatic necrosis (figure 3). In most cases, a combination of parenchymal and peripancreatic necrosis is seen.⁵ In the first 3–4 days of acute pancreatitis, CECT is unreliable for determining the extent of necrosis and the presence of collections.^{75 76} Only patients suspected of having abdominal catastrophes, such as perforation, bleeding or ischaemia, should have an urgent CECT.^{76 77} If patients fail to improve after 5–7 days of initial treatment, a CECT can determine the presence and extent of necrosis and (peri)pancreatic collections.^{4 64}

The terminology of (peri)pancreatic collections has changed in the revised Atlanta classification.⁵ In case of interstitial pancreatitis, these collections are referred to as 'acute pancreatic fluid collections', and in case of necrotising pancreatitis as 'acute necrotic collections' (ANC). Over time, acute collections either resolve spontaneously or mature with encapsulation of fluid and/or necrotic tissue. In ANC, this leads to walled-off necrosis




Figure 3 Peripancreatic necrosis—a form of necrotising pancreatitis.

(WON) (figure 4).^{5 78} The process of encapsulation mostly takes around 4–6 weeks. If the necrotic collections remain sterile, patients can be treated conservatively. Intervention is only indicated in case of infection. In sterile collections, interventions should only be considered in case of mechanical obstruction and/or failure to thrive which persists for 6–8 weeks.

INFECTED NECROTISING PANCREATITIS

Collections with necrosis (ie, ANC, WON) become infected in about one-third of patients.⁷⁹ Infected necrotising pancreatitis has a mortality of 15%.³ Infection can be diagnosed in three ways: (1) by gas configurations in the necrotic collection on imaging (figure 5),⁵ (2) by a positive gram stain or culture from a (percutaneous) fine-needle aspiration of the necrotic collection⁸⁰ or (3) suspected by clinical diagnosis. Clinical suspicion of infection is based on signs of infection (temperature >38.5°C, rising serum inflammatory markers) or when new/persistent organ failure occurs, which is typically most reliable after the initial phase of SIRS.^{4 81 82}




Figure 4 Walled-off necrosis—a form of necrotising pancreatitis.




Figure 5 Infected necrotic collection, with gas configurations, on day 20 after onset of disease. Not fully encapsulated.

If infection of a necrotic collection is proven or clinically highly suspected, antibiotic treatment is indicated. The preferred antibiotics are broad-spectrum, with the capability of penetrating the necrotic pancreatic tissue, according to local antibiotics protocol. A positive culture may be used to switch to targeted antibiotics. Although there are some series with high success rates of solely antibiotic treatment, most patients will eventually need an intervention (ie, catheter drainage, necrosectomy) to treat infected necrosis (figure 6).^{82–84} Antibiotics are therefore mostly used to support patients until collections become encapsulated (WON). This is presumed to facilitate safer interventions with a lower risk of bleeding and less reinterventions. Some experts, however, question this delay in treatment and suggest immediate catheter drainage.⁸⁵ A multicentre RCT* is currently comparing immediate versus postponed drainage in 104 patients with infected necrotising pancreatitis (POINTER trial; ISCRTN33682933).

Intervention

Historically, patients with necrotising pancreatitis underwent early laparotomy to debride the necrotic tissue. This was associated with high mortality rates, probably because these often severely ill patients could not endure the extra ‘hit’ of the surgical trauma.^{86 87} Current guidelines therefore advocate to delay interventions until the stage of WON.^{4 64} In a multicentre RCT*, 88 patients with infected necrotising pancreatitis were randomised between necrosectomy via laparotomy or a step-up approach. This approach consisted of percutaneous catheter drainage followed, in case of lack of clinical improvement, by video-assisted retroperitoneal debridement.⁸² The step-up approach reduced the composite end point (ie, death or major complications) from 69% to 40% (risk ratio (RR) 0.57; 95% CI 0.38 to 0.87; $p=0.006$). Another finding was that 35% of patients in the step-up arm could be treated with catheter drainage only, and did not require a necrosectomy.⁸²

The step-up approach is now considered the standard practice of care for patients with infected pancreatic necrosis and has been implemented in all major guidelines.^{4 64} Catheter drainage is only followed by necrosectomy when clinically indicated. Several methods of surgical necrosectomy are available, either open or minimally invasive techniques. As RCTs comparing these methods ‘head-to-head’ are lacking, the optimal method of necrosectomy




Figure 6 Flowchart acute pancreatitis. CECT, contrast-enhanced CT; ERC, endoscopic retrograde cholangiography; ES, endoscopic sphincterotomy; VARD, video-assisted retroperitoneal debridement.

remains unclear. Retrospective studies suggest a decreased risk of complications using minimally invasive techniques.^{88 89}

The step-up approach can also be performed endoscopically. In 2000, endosonography-guided transgastric necrosectomy was first described. It avoids general anaesthesia and may further reduce the surgical stress and complications.⁹⁰ The technique has since been reported in several retrospective cohort studies.^{91 92} A multicentre pilot RCT* in 20 patients found a reduction in the primary end point of proinflammatory response and in the combined secondary end point of major complications in patients undergoing endoscopic necrosectomy, compared with minimal invasive surgical necrosectomy.⁹³ Although promising, this was a pilot RCT and did not include a step-up approach. A subsequent multicentre RCT*, including 98 patients, comparing an endoscopic step-up approach with a surgical step-up approach has recently been completed and results are awaited (TENSION trial, ISRCTN09186711).⁸¹

AFTERCARE

Prevention of recurrence

Some 17%–22% of patients will have a recurrent pancreatitis and 8%–16% of patients will develop chronic pancreatitis.^{94–96} Several studies have attempted to reduce recurrence rates, most of them addressing the underlying cause of the disease. In patients with alcoholic pancreatitis, supervised alcohol abstinence should be advised, as continuation of alcohol consumption increases the risk of recurrence and ultimately of chronic pancreatitis.^{97–99} Smoking is an additional, but poorly recognised, risk factor for recurrent acute and chronic pancreatitis.⁹⁴ One RCT in 120 patients showed that repeated outpatient clinic visits with an intervention against alcohol consumption reduces the recurrence of pancreatitis, compared with a single intervention.¹⁰⁰ However, adherence to these abstinence programmes remains poor.¹⁰¹

In patients with biliary pancreatitis, cholecystectomy will reduce the risk of recurrence but its optimal timing has been debated. In severe biliary pancreatitis, it is common practice to delay cholecystectomy until the patient has recovered and local signs of inflammation have resolved or until at least 6 weeks after discharge.¹⁰² In mild biliary pancreatitis, current guidelines recommend cholecystectomy during the same hospital admission. Several clinical audits have shown that guideline adherence is poor, cholecystectomy is often delayed.^{99 103–107} Concerns about the perceived increased difficulty of surgical dissection after pancreatitis, resulting in higher surgical complication rates,^{108 109} and logistical challenges with busy emergency theatre lists^{104 110} have probably contributed to these delays. A recent multicentre RCT* in 266 patients with mild biliary pancreatitis demonstrated that same-admission cholecystectomy reduced the recurrence rate of gallstone-related complications from 17% to 5% as compared with interval cholecystectomy after 4 weeks (RR 0.28, 95% CI 0.12 to 0.66; $p=0.002$).¹¹¹ This included a reduction of recurrent biliary pancreatitis from 9% to 2% (RR 0.27; 95% CI 0.08 to 0.92, $p=0.03$). Same-admission cholecystectomy was not associated with increased technical difficulty or complications and decreased overall costs.^{111 112}

The value of cholecystectomy has also been investigated in acute idiopathic pancreatitis. A multicentre RCT in 85 patients showed that routine elective laparoscopic cholecystectomy reduced the rate of recurrent pancreatitis from 30% to 10% ($p=0.016$) as compared with conservative management. However, since routine EUS was not included in the work-up of (presumed) idiopathic pancreatitis, the treatment effect of cholecystectomy may have been overestimated in this trial. Many patients with presumed

idiopathic pancreatitis patients may in fact have had acute biliary pancreatitis. This was also reflected in the large percentage (59%) of gallbladders in which biliary stones or sludge were found at pathological examination in this RCT.¹¹³

Exocrine and endocrine insufficiency

Due to extensive loss of pancreatic tissue in necrotising pancreatitis, there is a reduction of both endocrine and/or exocrine pancreatic function in 19%–80% of all patients.^{114–117} Awareness on these conditions may support timely treatment in order to prevent complications from diabetes or malnutrition from malabsorption.

Future

This review summarised the best available evidence in acute pancreatitis, mostly based on RCTs. Although many clinical questions have been addressed by RCTs, many questions remain, for instance regarding fluid and pain management but also on the use and timing of interventions such as ERC and the ‘step-up approach’. Besides these important questions, no single RCT has reported on an effective treatment to halt the early sequence of severe systemic inflammation, ultimately leading to multiple organ failure and death in acute pancreatitis. Further innovative studies and RCTs are needed to find such a treatment and to improve outcomes in acute pancreatitis.

Acknowledgements Several randomised controlled trials(*) described in this review were performed by the *Dutch Pancreatitis Study Group*.

Contributors Conception or design of the work: SMvD, NDLH, Hvs, PF, HvG, MJB, MGB. Data collection: SMvD, NDLH, Hvs, MJB, MGB. Data analysis and interpretation: SMvD, NDLH, Hvs, PF, HvG, MJB, MGB. Drafting the article: SMvD, NDLH, MJB, MGB. Critical revision of the article: SMvD, NDLH, Hvs, PF, HvG, MJB, MGB. Final approval of the version to be published: SMvD, NDLH, Hvs, PF, HvG, MJB, MGB.

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Hospital Episode Statistics. *Admitted Patient Care - England, 2014-15 (NS)*. (accessed 25 Nov 2015).
- Roberts SE, Morrison-Rees S, John A, *et al*. The incidence and aetiology of acute pancreatitis across Europe. *Pancreatology* 2017;17:155–65.
- Banks PA, Freeman ML. Practice Parameters Committee of the American College of Gastroenterology. practice guidelines in acute pancreatitis. *Am J Gastroenterol* 2006;101:2379–400.
- Working Group IAP/APA acute pancreatitis guidelines. IAP/APA evidence-based guidelines for the management of acute pancreatitis. *Pancreatology* 2013;13:e1–15.
- Banks PA, Bollen TL, Dervenis C, *et al*. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013;62:102–11.
- Frey CF, Zhou H, Harvey DJ, *et al*. The incidence and case-fatality rates of acute biliary, alcoholic, and idiopathic pancreatitis in California, 1994–2001. *Pancreas* 2006;33:336–44.
- Yadav D, Lowenfels AB. Trends in the epidemiology of the first attack of acute pancreatitis: a systematic review. *Pancreas* 2006;33:323–30.
- Nesvaderani M, Eslick GD, Vagg D, *et al*. Epidemiology, aetiology and outcomes of acute pancreatitis: a retrospective cohort study. *Int J Surg* 2015;23:68–74.
- Smith I, Ramesh J, Kyanam Kabir Baig KR, *et al*. Emerging role of endoscopic ultrasound in the Diagnostic evaluation of idiopathic pancreatitis. *Am J Med Sci* 2015;350:229–34.
- Mounzer R, Langmead CJ, Wu BU, *et al*. Comparison of existing clinical scoring systems to predict persistent organ failure in patients with acute pancreatitis. *Gastroenterology* 2012;142:1476–82.
- Perel P, Roberts I, Ker K. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev* 2013:CD000567.
- Wu BU, Hwang JQ, Gardner TH, *et al*. Lactated Ringer's solution reduces systemic inflammation compared with saline in patients with acute pancreatitis. *Clin Gastroenterol Hepatol* 2011;9:710–7.

- 13 Young P, Bailey M, Beasley R, *et al.* Effect of a Buffered Crystalloid solution vs saline on acute kidney injury among Patients in the Intensive Care Unit: the SPLIT Randomized clinical trial. *JAMA* 2015;314:314.
- 14 Shaw AD, Schermer CR, Lobo DN, *et al.* Impact of intravenous fluid composition on outcomes in patients with systemic inflammatory response syndrome. *Crit Care* 2015;19:334.
- 15 Raghunathan K, Shaw A, Nathanson B, *et al.* Association between the choice of IV crystalloid and in-hospital mortality among critically ill adults with Sepsis. *Crit Care Med* 2014;42:1585–91.
- 16 Yunos NM, Bellomo R, Hegarty C, *et al.* Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA* 2012;308:1566–72.
- 17 Mao EQ, Tang YQ, Fei J, *et al.* Fluid therapy for severe acute pancreatitis in acute response stage. *Chin Med J* 2009;122:169–73.
- 18 Mao EQ, Fei J, Peng YB, *et al.* Rapid hemodilution is associated with increased Sepsis and mortality among patients with severe acute pancreatitis. *Chin Med J* 2010;123:1639–44.
- 19 Buxbaum JL, Quezada M, Da B, *et al.* Early Aggressive Hydration hastens clinical improvement in Mild acute pancreatitis. *Am J Gastroenterol* 2017;112:797–803.
- 20 Wang MD, Ji Y, Xu J, *et al.* Early goal-directed fluid therapy with fresh frozen plasma reduces severe acute pancreatitis mortality in the intensive care unit. *Chin Med J* 2013;126:1987–8.
- 21 Wilms B, Meffert K, Schultes B. Procaininfusion zur Schmerzbehandlung bei akuter pankreatitis. *DMW - Deutsche Medizinische Wochenschrift* 2010;135:2290–5.
- 22 Lauer P, Bronisch HJ, Henniges UM, *et al.* Effects of systemic administration of a local anesthetic on pain in acute pancreatitis: a randomized clinical trial. *Pancreas* 2011;40:673–9.
- 23 Jakobs R, Adamek MU, von Bubnoff AC, *et al.* Buprenorphine or procaine for pain relief in acute pancreatitis. A prospective randomized study. *Scand J Gastroenterol* 2000;35:1319–23.
- 24 Kahl S, Zimmermann S, Pross M, *et al.* Procaine hydrochloride fails to relieve pain in patients with acute pancreatitis. *Digestion* 2004;69:5–9.
- 25 Blamey SL, Finlay IG, Carter DC, *et al.* Analgesia in acute pancreatitis: comparison of buprenorphine and pethidine. *Br Med J* 1984;288:1494–5.
- 26 Stevens M, Esler R, Asher G. Transdermal fentanyl for the management of acute pancreatitis pain. *Appl Nurs Res* 2002;15:102–10.
- 27 Peiró AM, Martínez J, Martínez E, *et al.* Efficacy and tolerance of metamizole versus morphine for acute pancreatitis pain. *Pancreatol* 2008;8:25–9.
- 28 Gülen B, Dur A, Serinken M, *et al.* Pain treatment in patients with acute pancreatitis: a randomized controlled trial. *Turk J Gastroenterol* 2016;27:192–6.
- 29 Ona B X, Rigau Comas D, Urrutia G. Opioids for acute pancreatitis pain. Ona B X, ed. *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd, 2013.
- 30 Meng W, Yuan J, Zhang C, *et al.* Parenteral analgesics for pain relief in acute pancreatitis: a systematic review. *Pancreatol* 2013;13:201–6.
- 31 Bradley EL. A clinically based classification system for acute pancreatitis. Summary of the International Symposium on acute pancreatitis, Atlanta, Ga, September 11 through 13, 1992. *Arch Surg* 1993;128:586–90.
- 32 Ammori BJ, Leeder PC, King RF, *et al.* Early increase in intestinal permeability in patients with severe acute pancreatitis: correlation with endotoxemia, organ failure, and mortality. *J Gastrointest Surg* 1999;3:252–62.
- 33 García-Barrasa A, Borobia FG, Pallares R, *et al.* A double-blind, placebo-controlled trial of ciprofloxacin prophylaxis in patients with acute necrotizing pancreatitis. *J Gastrointest Surg* 2009;13:768–74.
- 34 Isenmann R, Rünzi M, Kron M, *et al.* Prophylactic antibiotic treatment in patients with predicted severe acute pancreatitis: a placebo-controlled, double-blind trial. *Gastroenterology* 2004;126:997–1004.
- 35 Dellinger EP, Tellado JM, Soto NE, *et al.* Early antibiotic treatment for severe acute necrotizing pancreatitis. *Ann Surg* 2007;245:674–83.
- 36 Lim CL, Lee W, Liew YX, *et al.* Role of antibiotic prophylaxis in necrotizing pancreatitis: a meta-analysis. *J Gastrointest Surg* 2015;19:480–91.
- 37 Villatoro E, Mulla M, Larvin M. Antibiotic therapy for prophylaxis against infection of pancreatic necrosis in acute pancreatitis. In: Larvin M, ed. *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd, 2010.
- 38 Oláh A, Belágyi T, Póto L, *et al.* Symbiotic control of inflammation and infection in severe acute pancreatitis: a prospective, randomized, double blind study. *Hepatogastroenterology* 2007;54:590–4.
- 39 Oláh A, Belágyi T, Issekutz A, *et al.* Randomized clinical trial of specific Lactobacillus and fibre supplement to early enteral nutrition in patients with acute pancreatitis. *Br J Surg* 2002;89:1103–7.
- 40 Besselink MG, van Santvoort HC, Buskens E, *et al.* Probiotic prophylaxis in predicted severe acute pancreatitis: a randomised, double-blind, placebo-controlled trial. *Lancet* 2008;371:651–9.
- 41 Van Felius ID, Akkermans LM, Bosscha K, *et al.* Interdigestive small bowel motility and duodenal bacterial overgrowth in experimental acute pancreatitis. *Neurogastroenterol Motil* 2003;15:267–76.
- 42 Rahman SH, Ammori BJ, Holmfeld J, *et al.* Intestinal Hypoperfusion contributes to gut barrier failure in severe acute pancreatitis. *J Gastrointest Surg* 2003;7:26–36.
- 43 Nieuwenhuijs VB, Verheem A, van Duijvenbode-Beumer H, *et al.* The role of interdigestive small bowel motility in the regulation of gut microflora, bacterial overgrowth, and bacterial translocation in rats. *Ann Surg* 1998;228:188–93.
- 44 Fritz S, Hackert T, Hartwig W, *et al.* Bacterial translocation and infected pancreatic necrosis in acute necrotizing pancreatitis derives from small bowel rather than from Colon. *Am J Surg* 2010;200:111–7.
- 45 Besselink MG, van Santvoort HC, Renooij W, *et al.* Intestinal barrier dysfunction in a randomized trial of a specific probiotic composition in acute pancreatitis. *Ann Surg* 2009;250:712–9.
- 46 Marik PE. What is the best way to feed patients with pancreatitis? *Curr Opin Crit Care* 2009;15:131–8.
- 47 McClave SA, Heyland DK. The physiologic response and associated clinical benefits from provision of early enteral nutrition. *Nutr Clin Pract* 2009;24:305–15.
- 48 Al-Omran M, Albalawi ZH, Tashkandi MF, *et al.* Enteral versus parenteral nutrition for acute pancreatitis. *Cochrane Database Syst Rev* 2010:CD002837.
- 49 Bakker OJ, van Brunschot S, Farre A, *et al.* Timing of enteral nutrition in acute pancreatitis: meta-analysis of individuals using a single-arm of randomised trials. *Pancreatol* 2014;14:340–6.
- 50 Wereszczynska-Siemiatkowska U, Swidnicka-Siergiejko A, Siemiatkowski A, *et al.* Early enteral nutrition is superior to delayed enteral nutrition for the prevention of infected necrosis and mortality in acute pancreatitis. *Pancreas* 2013;42:640–6.
- 51 Sun JK, Mu XW, Li WQ, *et al.* Effects of early enteral nutrition on immune function of severe acute pancreatitis patients. *World J Gastroenterol* 2013;19:917–22.
- 52 Bakker OJ, van Brunschot S, van Santvoort HC, *et al.* Early versus on-demand nasoenteric tube feeding in acute pancreatitis. *N Engl J Med* 2014;371:1983–93.
- 53 Stimac D, Poropat G, Hauser G, *et al.* Early nasojejunal tube feeding versus nil-by-mouth in acute pancreatitis: a randomized clinical trial. *Pancreatol* 2016;16:523–8.
- 54 Eatock FC, Chong P, Menezes N, *et al.* A randomized study of early nasogastric versus nasojejunal feeding in severe acute pancreatitis. *Am J Gastroenterol* 2005;100:432–9.
- 55 Kumar A, Singh N, Prakash S, *et al.* Early enteral nutrition in severe acute pancreatitis: a prospective randomized controlled trial comparing nasojejunal and nasogastric routes. *J Clin Gastroenterol* 2006;40:431–4.
- 56 Singh N, Sharma B, Sharma M, *et al.* Evaluation of Early Enteral Feeding through Nasogastric and Nasojejunal Tube in Severe Acute pancreatitis. *Pancreas* 2012;41:153–9.
- 57 Chang YS, Fu HQ, Xiao YM, *et al.* Nasogastric or nasojejunal feeding in predicted severe acute pancreatitis: a meta-analysis. *Crit Care* 2013;17:R118.
- 58 Eckerwall GE, Tingstedt BB, Bergenzaun PE, *et al.* Immediate oral feeding in patients with mild acute pancreatitis is safe and may accelerate recovery – a randomized clinical study. *Clin Nutr* 2007;26:758–63.
- 59 Moraes JM, Felga GE, Chebli LA, *et al.* A full solid diet as the initial meal in mild acute pancreatitis is safe and result in a shorter length of hospitalization: results from a prospective, randomized, controlled, double-blind clinical trial. *J Clin Gastroenterol* 2010;44:517–22.
- 60 Teich N, Aghdassi A, Fischer J, *et al.* Optimal timing of oral refeeding in mild acute pancreatitis: results of an open randomized multicenter trial. *Pancreas* 2010;39:1088–92.
- 61 Oriá A, Cimmino D, Ocampo C, *et al.* Early endoscopic intervention versus early conservative management in patients with acute gallstone pancreatitis and biliopancreatic obstruction: a randomized clinical trial. *Ann Surg* 2007;245:10–17.
- 62 Fölsch UR, Nitsche R, Lütke R, *et al.* Early ERCP and papillotomy compared with conservative treatment for acute biliary pancreatitis. the german Study Group on acute biliary pancreatitis. *N Engl J Med* 1997;336:237–42.
- 63 Nowak A, Marek TA, Nowakowska-Dulawa E, *et al.* Biliary pancreatitis needs endoscopic retrograde cholangiopancreatography with endoscopic sphincterotomy for cure. *Endoscopy* 1998;30:A 256–A 259.
- 64 Tenner S, Baillye J, DeWitt J, *et al.* American College of Gastroenterology Guideline: Management of Acute Pancreatitis. *Am J Gastroenterol* 2013;108:1400–15; 1416.
- 65 Neoptolemos JP, Carr-Locke DL, London NJ, *et al.* Controlled trial of urgent endoscopic retrograde cholangiopancreatography and endoscopic sphincterotomy versus conservative treatment for acute pancreatitis due to gallstones. *Lancet* 1988;2:979–83.
- 66 Kiriya S, Takada T, Strasberg SM, *et al.* TG13 guidelines for diagnosis and severity grading of acute cholangitis (with videos). *J Hepatobiliary Pancreat Sci* 2013;20:24–34.
- 67 Fan ST, Lai EC, Mok FP, *et al.* Early treatment of acute biliary pancreatitis by endoscopic papillotomy. *N Engl J Med* 1993;328:228–32.
- 68 Acosta JM, Katkhouda N, Deban KA, *et al.* Early ductal decompression versus conservative management for gallstone pancreatitis with ampullary obstruction: a prospective randomized clinical trial. *Ann Surg* 2006;243:33–40.
- 69 van Geenen EJ, van Santvoort HC, Besselink MG, *et al.* Lack of consensus on the role of endoscopic retrograde cholangiopancreatography in acute biliary pancreatitis in published meta-analyses and guidelines: a systematic review. *Pancreas* 2013;42:774–80.
- 70 Schepers NJ, Bakker OJ, Besselink MG, *et al.* Early biliary decompression versus conservative treatment in acute biliary pancreatitis (APEC trial): study protocol for a randomized controlled trial. *Trials* 2016;17:5.

Recent advances in clinical practice

- 71 Liu CL, Fan ST, Lo CM, *et al.* Comparison of early endoscopic ultrasonography and endoscopic retrograde cholangiopancreatography in the management of acute biliary pancreatitis: a prospective randomized study. *Clin Gastroenterol Hepatol* 2005;3:1238–44.
- 72 De Lisi S, Leandro G, Buscarini E. Endoscopic ultrasonography versus endoscopic retrograde cholangiopancreatography in acute biliary pancreatitis: a systematic review. *Eur J Gastroenterol Hepatol* 2011;23:367–74.
- 73 Anderloni A, Galeazzi M, Ballarè M, *et al.* Early endoscopic ultrasonography in acute biliary pancreatitis: a prospective pilot study. *World J Gastroenterol* 2015;21:10427–34.
- 74 Romagnuolo J, Currie G; Calgary Advanced Therapeutic Endoscopy Center study group. Noninvasive vs. selective invasive biliary imaging for acute biliary pancreatitis: an economic evaluation by using decision tree analysis. *Gastrointest Endosc* 2005;61:86–97.
- 75 Spanier BW, Nio Y, van der Hulst RW, *et al.* Practice and yield of early CT scan in acute pancreatitis: a dutch Observational Multicenter Study. *Pancreatology* 2010;10:222–8.
- 76 Bollen TL, Singh VK, Maurer R, *et al.* A comparative evaluation of radiologic and clinical scoring systems in the early prediction of severity in acute pancreatitis. *Am J Gastroenterol* 2012;107:612–9.
- 77 Bollen TL. Acute pancreatitis: international classification and nomenclature. *Clin Radiol* 2016;71:121–33.
- 78 Sarathi Patra P, Das K, Bhattacharyya A, *et al.* Natural resolution or intervention for fluid collections in acute severe pancreatitis. *Br J Surg* 2014;101:1721–8.
- 79 van Santvoort HC, Bakker OJ, Bollen TL, *et al.* A conservative and minimally invasive approach to necrotizing pancreatitis improves outcome. *Gastroenterology* 2011;141:1254–63.
- 80 van Baal MC, Bollen TL, Bakker OJ, *et al.* The role of routine fine-needle aspiration in the diagnosis of infected necrotizing pancreatitis. *Surgery* 2014;155:442–8.
- 81 van Brunschot S, van Grinsven J, Voermans RP, *et al.* Transluminal endoscopic step-up approach versus minimally invasive surgical step-up approach in patients with infected necrotising pancreatitis (TENSION trial): design and rationale of a randomised controlled multicenter trial [ISRCTN09186711]. *BMC Gastroenterol* 2013;13:161.
- 82 van Santvoort HC, Besselink MG, Bakker OJ, *et al.* A step-up approach or open necrosectomy for necrotizing pancreatitis. *N Engl J Med* 2010;362:1491–502.
- 83 Adler DG, Chari ST, Dahl TJ, *et al.* Conservative management of infected necrosis complicating severe acute pancreatitis. *Am J Gastroenterol* 2003;98:98–103.
- 84 Runzi M, Niebel W, Goebell H, *et al.* Severe acute pancreatitis: nonsurgical treatment of infected necroses. *Pancreas* 2005;30:195–9.
- 85 van Grinsven J, van Brunschot S, Bakker OJ, *et al.* Diagnostic strategy and timing of intervention in infected necrotizing pancreatitis: an international expert survey and case vignette study. *HPB* 2016;18:49–56.
- 86 Besselink MG, Verwer TJ, Schoenmaeckers EJ, *et al.* Timing of surgical intervention in necrotizing pancreatitis. *Arch Surg* 2007;142:1194–201.
- 87 Fernández-del Castillo C, Rattner DW, Makary MA, *et al.* Débridement and closed packing for the treatment of necrotizing pancreatitis. *Ann Surg* 1998;228:676–84.
- 88 Raraty MGT, Halloran CM, Dodd S, *et al.* Minimal access retroperitoneal pancreatic necrosectomy. *Ann Surg* 2010;251:787–93.
- 89 Gomatos IP, Halloran CM, Ghaneh P, *et al.* Outcomes from minimal access retroperitoneal and open pancreatic necrosectomy in 394 patients with necrotizing pancreatitis. *Ann Surg* 2016;263:992–1001.
- 90 Seifert H, Wehrmann T, Schmitt T, *et al.* Retroperitoneal endoscopic debridement for infected peripancreatic necrosis. *Lancet* 2000;356:653–5.
- 91 van Brunschot S, Fockens P, Bakker OJ, *et al.* Endoscopic transluminal necrosectomy in necrotising pancreatitis: a systematic review. *Surg Endosc* 2014;28:1425–38.
- 92 Haghshenas Kashani A, Laurence JM, Kwan V, *et al.* Endoscopic necrosectomy of pancreatic necrosis: a systematic review. *Surg Endosc* 2011;25:3724–30.
- 93 Bakker OJ, van Santvoort HC, van Brunschot S, *et al.* Endoscopic transgastric vs surgical necrosectomy for infected necrotizing pancreatitis: a randomized trial. *JAMA* 2012;307:1053–61.
- 94 Ahmed Ali U, Issa Y, Hagensaers JC, *et al.* Risk of recurrent pancreatitis and progression to chronic pancreatitis after a first episode of acute pancreatitis. *Clin Gastroenterol Hepatol* 2016;14:738–46.
- 95 Yadav D, O'Connell M, Papachristou GI. Natural history following the first attack of acute pancreatitis. *Am J Gastroenterol* 2012;107:1096–103.
- 96 Lankisch PG, Breuer N, Bruns A, *et al.* Natural history of acute pancreatitis: a long-term population-based study. *Am J Gastroenterol* 2009;104:2797–805.
- 97 Sand J, Lankisch PG, Nordback I. Alcohol consumption in patients with acute or chronic pancreatitis. *Pancreatology* 2007;7:147–56.
- 98 Pelli H, Lappalainen-Lehto R, Piironen A, *et al.* Risk factors for recurrent acute alcohol-associated pancreatitis: a prospective analysis. *Scand J Gastroenterol* 2008;43:614–21.
- 99 NCEPOD - Acute pancreatitis: treat the cause (2016). *Natl. Confid. Enq. into Patient Outcome Death* 2016.
- 100 Nordback I, Pelli H, Lappalainen-Lehto R, *et al.* The recurrence of acute alcohol-associated pancreatitis can be reduced: a randomized controlled trial. *Gastroenterology* 2009;136:848–55.
- 101 Singer MV, Pfützer RH, Kiefer F. Striving for abstinence in alcoholic pancreatitis: act of humanity, economic necessity, or flogging a dead horse after all? *Gastroenterology* 2009;136:757–60.
- 102 Nealon WH, Bawduniak J, Walser EM. Appropriate timing of cholecystectomy in patients who present with moderate to severe gallstone-associated acute pancreatitis with peripancreatic fluid collections. *Ann Surg* 2004;239:741–9; discussion 749–51.
- 103 Pezzilli R, Uomo G, Gabbriellini A, *et al.* A prospective multicentre survey on the treatment of acute pancreatitis in Italy. *Dig Liver Dis* 2007;39:838–46.
- 104 Lankisch PG, Weber-Dany B, Lerch MM. Clinical perspectives in pancreatology: compliance with acute pancreatitis guidelines in Germany. *Pancreatology* 2005;5:591–3.
- 105 Sargen K, Kingsnorth AN. Management of gallstone pancreatitis: effects of deviation from clinical guidelines. *JOP* 2001;2:317–22.
- 106 Monkhouse SJ, Court EL, Dash I, *et al.* Two-week target for laparoscopic cholecystectomy following gallstone pancreatitis is achievable and cost neutral. *Br J Surg* 2009;96:751–5.
- 107 Cameron DR, Goodman AJ. Delayed cholecystectomy for gallstone pancreatitis: re-admissions and outcomes. *Ann R Coll Surg Engl* 2004;86:358–62.
- 108 Kelly TR, Wagner DS. Gallstone pancreatitis: a prospective randomized trial of the timing of surgery. *Surgery* 1988;104:600–5.
- 109 Johnstone M, Marriott P, Royle TJ, *et al.* The impact of timing of cholecystectomy following gallstone pancreatitis. *Surgeon* 2014;12:134–40.
- 110 Nguyen GC, Boudreau H, Jagannath SB. Hospital volume as a predictor for undergoing cholecystectomy after admission for acute biliary pancreatitis. *Pancreas* 2010;39:e42–7.
- 111 da Costa DW, Bouwense SA, Schepers NJ, *et al.* Same-admission versus interval cholecystectomy for mild gallstone pancreatitis (PONCHO): a multicentre randomised controlled trial. *Lancet* 2015;386:1261–8.
- 112 da Costa DW, Dijkstra LM, Bouwense SA, *et al.* Cost-effectiveness of same-admission versus interval cholecystectomy after mild gallstone pancreatitis in the PONCHO trial. *Br J Surg* 2016;103:1695–703.
- 113 Rätty S, Pulkkinen J, Nordback I, *et al.* Can Laparoscopic Cholecystectomy prevent recurrent idiopathic acute pancreatitis? *Ann Surg* 2015;262:736–41.
- 114 Rana SS, Bhasin DK, Rao C, *et al.* Comparative evaluation of structural and functional changes in pancreas after endoscopic and surgical management of pancreatic necrosis. *Ann Gastroenterol* 2014;27:162–6.
- 115 Gupta R, Wig JD, Bhasin DK, *et al.* Severe acute pancreatitis: the life after. *J Gastrointest Surg* 2009;13:1328–36.
- 116 Reddy MS, Singh S, Singh R, *et al.* Morphological and functional outcome after pancreatic necrosectomy and lesser sac lavage for necrotizing pancreatitis. *Indian J Gastroenterol* 2007;26:217–20.
- 117 Chandrasekaran P, Gupta R, Shenvi S, *et al.* Prospective comparison of long term outcomes in patients with severe acute pancreatitis managed by operative and non operative measures. *Pancreatology* 2015;15:478–84.



Acute pancreatitis: recent advances through randomised trials

Sven M van Dijk, Nora D L Hallensleben, Hjalmar C van Santvoort, Paul Fockens, Harry van Goor, Marco J Bruno and Marc G Besselink

Gut 2017 66: 2024-2032 originally published online August 24, 2017
doi: 10.1136/gutjnl-2016-313595

Updated information and services can be found at:
<http://gut.bmj.com/content/66/11/2024>

These include:

References

This article cites 111 articles, 2 of which you can access for free at:
<http://gut.bmj.com/content/66/11/2024#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[GUT Recent advances in clinical practice](#) (100)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>