

Minimally invasive versus open lateral pancreaticojejunostomy in patients with painful chronic pancreatitis: systematic review

Roberto M. Montorsi^{1,2,3,*} , Michiel F.G. Francken^{1,2,4}, Marja A. Boermeester^{1,2} , Olivier R. Busch^{1,2} , Freek Daams^{2,5}, Thilo Hackert⁶ , Roel Haen⁷, Markus W. Hollmann⁸, Hjalmar C. van Santvoort^{9,10}, and Marc G. Besselink^{1,2} ; for the Dutch Pancreatitis Study Group

¹Department of Surgery, Amsterdam UMC, location University of Amsterdam, Amsterdam, The Netherlands

²Department of Surgery, Amsterdam Gastroenterology Endocrinology Metabolism, Amsterdam, The Netherlands

³Department of General and Pancreatic Surgery, The Pancreas Institute, University of Verona Hospital Trust, Verona, Italy

⁴Department of Research and Development, St Antonius Hospital, Nieuwegein, The Netherlands

⁵Department of Surgery, Amsterdam UMC, location Vrije Universiteit, Amsterdam, The Netherlands

⁶Department of General, Visceral and Thoracic Surgery, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

⁷Department of Surgery, Erasmus University Medical Center, Rotterdam, The Netherlands

⁸Department of Anesthesiology, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

⁹Department of Surgery, St. Antonius Hospital, Nieuwegein, The Netherlands

¹⁰Department of Surgery, University Medical Center Utrecht, Utrecht, The Netherlands

*Correspondence to: Roberto Maria Montorsi, MD, PhD, Department of Surgery, Amsterdam Gastroenterology Endocrinology Metabolism, University of Amsterdam, De Boelelaan 1117 (ZH-7F), Amsterdam 1081 HV, The Netherlands (E-mail: montorsi.imd@gmail.it)

Abstract

Background: Patients with painful chronic pancreatitis combined with a dilated main pancreatic duct and a normal size pancreatic head are treated according to guidelines by lateral pancreaticojejunostomy (LPJ). This systematic review compared outcomes of minimally invasive LPJ and open LPJ.

Methods: From 1 January 2000 until 13 November 2023, series reporting on minimally invasive LPJ and open LPJ in patients with symptomatic chronic pancreatitis were included. This study was structured in accordance with the PRISMA guidelines. The primary outcome was intraoperative and postoperative complications. Secondary outcomes included long-term clinical outcomes.

Results: Overall, 19 retrospective studies were included. Morbidity rate ranged from 0% to 57% after minimally invasive LPJ versus 4% to 68% after open LPJ (median: 25, i.q.r.: 23). Length of hospital stay ranged from 5 to 7 days after minimally invasive LPJ and from 6 to 16 days after open LPJ. The rate of pain relief ranged from 62% to 91% after open LPJ (median: 78.5, i.q.r.: 23) and from 71% to 100% (median: 82.5, i.q.r.: 12.5) after minimally invasive LPJ respectively. New-onset endocrine insufficiency ranged from 21% to 22% in minimally invasive LPJ and 19% to 26% after open LPJ. New-onset exocrine insufficiency was shown in 11% to 27% in minimally invasive LPJ versus 8% to 26% after open LPJ. Weight gain ranged from 60% to 100% (median: 97, i.q.r.: 23) after minimally invasive LPJ.

Discussion: This systematic review suggested that minimally invasive LPJ can be performed safely in selected patients with symptomatic chronic pancreatitis. Phase 2 randomized trials should assess potential short-term benefits such as postoperative pain and length of hospital stay after minimally invasive LPJ.

Introduction

Chronic pancreatitis (CP) can be characterized by pain, endocrine and exocrine pancreatic insufficiency, and increased risk of pancreatic cancer, resulting in reduced quality of life^{1,2}. Management of CP aims to increase patients' quality of life (QoL) by relieving symptoms and preventing disease progression. Chronic pain in CP is a multifactorial problem^{3,4}. Approximately 75% of patients with CP experience chronic pain and 30% of these patients need medical therapy and eventually require intervention⁵.

Several randomized studies have demonstrated that surgery is the most effective treatment, especially in patients with CP and a dilated main pancreatic duct^{6–8}. According to current guidelines^{9–12}, lateral pancreaticojejunostomy (LPJ) is the procedure

of choice in patients with chronic pain and is indicated in patients in whom the main pancreatic duct is dilated (>5 mm) but without an enlargement of the pancreatic head (≤40 mm). Surgical LPJ is associated with a low risk of mortality and a relatively low risk of postoperative complications⁹.

Following surgery, these patients are prone to a challenging pain therapy. Most patients are long-standing opioid users and may present with opioid misuse, abuse, tolerance, dependency, or even opioid-induced hyperalgesia, lowering sensitivity to opioids or, in the case of hyperalgesia, they might even suffer from increased pain following opioid administration. Therefore, minimally invasive surgery (MIS) LPJ has been suggested as a means of minimizing direct postoperative pain, thus enhancing time to functional recovery while reducing length of hospital

Received: April 29, 2024. Revised: September 11, 2024. Accepted: October 08, 2024

© The Author(s) 2025. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

stay, as long as the effectivity on CP-related pain is as good as open surgery. Large series and systematic reviews comparing MIS and open LPJ are currently lacking.

The aim of this systematic review was to report the surgical and long-term outcomes of MIS compared to open LPJ.

Methods

This systematic review was structured in accordance with the PRISMA¹³ guidelines. The study protocol was registered with the International Prospective Register of Systematic Reviews¹⁴ (PROSPERO) on 25 December 2023 (registration number CRD42023493554).

Information sources and search strategy

A systematic literature search was performed in PubMed, EMBASE (Ovid), and the Cochrane Library. The search included the words ("Pancreatitis"[Mesh] OR "Pancreatic Diseases"[Mesh] OR pancrea*[tiab]) AND ("Minimally Invasive Surgical Procedures"[Mesh] OR "Robotic Surgical Procedures"[Mesh] OR "Robotics"[Mesh] OR "Laparoscopy"[Mesh] OR "Pancreatitis/surgery"[Mesh:NoExp] OR "Pancreatitis, Chronic/surgery"[Mesh] OR robot*[tiab] OR RDP[tiab] OR minimally invasiv*[tiab] OR minimal invasiv*[tiab] OR mini-invasiv*[tiab] OR laparoscop*[tiab] OR open[tiab]) AND ("Pancreaticojejunostomy"[MeSH] OR pancreatojejunostom*[tiab] OR pancreaticojejunostom*[tiab] OR pancreato jejuna*[tiab] OR pancreatico jejuna*[tiab] OR puestow[tiab]) NOT (("Animals"[MeSH Terms] OR "models, animal"[MeSH Terms] OR "Animal Experimentation"[MeSH Terms] OR rat[tiab] OR rats[tiab] OR mice[tiab] OR mouse[tiab] OR murine[tiab] OR murines[tiab] OR rodent[tiab] OR rodents[tiab] OR rabbit[tiab] OR rabbits[tiab] OR cat[tiab] OR cats[tiab] OR dog[tiab] OR dogs[tiab] OR pig[tiab] OR pigs[tiab] OR cow[tiab] OR cows[tiab] OR monkey[tiab] OR monkeys[tiab] OR goat[tiab] OR goats[tiab] OR horse[tiab] OR horses[tiab] OR ape[tiab] OR apes[tiab] OR gorilla[tiab] OR gorillas[tiab] OR sheep[tiab] OR sheeps[tiab] OR ovine[tiab] OR lamb[tiab] OR swine[tiab] OR swines[tiab] OR porcine[tiab] OR pup[tiab] OR pups[tiab] OR canine[tiab] OR beagle[tiab]) NOT "Humans"[MeSH Terms]) NOT ("Letter" [Publication Type] OR "Comment" [Publication Type] OR "Editorial" [Publication Type] OR letter[ti] OR editorial[ti] OR comment*[ti]) AND ("2000/01/01"[Date—Publication] : "2023/11/13"[Date—Publication]). Duplicates were removed.

Eligibility criteria

All published studies reporting on MIS and/or open LPJ in patients with chronic pancreatitis were screened for eligibility. Both comparative and non-comparative studies were included. Randomized clinical trials (RCTs), prospective and retrospective observational studies were included (1 January 2000–13 November 2023). Case series reporting on less than five MIS LPJ, trial protocols, systematic reviews, meta-analyses, conference abstracts, secondary publications of previously published studies, letters and commentaries were excluded. Publications lacking full text and publications in languages other than English were also excluded. In order to prevent a much larger control group than intervention group, studies with <50 open LPJ were excluded.

Data collection process

Two authors (R.M.M. and M.F.G.F.) independently screened literature on title and abstract in a blinded fashion and ultimately on full text using the aforementioned eligibility

criteria using Rayyan¹⁵ (version 2022). Disagreements were resolved by consensus and, if necessary, by the opinion of a third reviewer (M.G.B.).

Data items

Data extraction included the publication details (for example study title, publication date, authors, study design), baseline characteristics (for example number of patients, sex, age, clinical characteristics, interventions (for example type of surgical approach, intraoperative variables) and postoperative characteristics (for example postoperative pancreatic fistula¹⁶, delayed gastric emptying¹⁷, postpancreatectomy haemorrhage¹⁸, bile leak¹⁹, chyle leak²⁰, postpancreatectomy acute pancreatitis²¹, general postoperative complications, Clavien–Dindo classification²², quality of life, long-term outcomes). Primary and secondary endpoints were extracted additionally.

Outcome

The primary outcomes considered intraoperative (for example operative time, estimated blood loss) and postoperative (for example morbidity rate, any postoperative complication regardless of Clavien–Dindo grade, length of hospital stay, readmission, mortality) data. Secondary outcomes included long-term outcomes (for example pain relief, endocrine insufficiency, exocrine insufficiency, weight gain, medical costs).

Risk of bias assessment

The quality of the included studies was assessed by two independent reviewers (R.M.M. and M.F.). The methodological index for non-randomized studies (MINORS)²³ and Risk of Bias 2 (RoB2)²⁴ criteria were used for non-randomized and randomized studies respectively. As stated by MINORS, the quality of comparative and non-comparative non-randomized studies was assessed with eight criteria. In case of comparative studies, the additional criteria assessed were the adequacy of the control group, contemporary groups, baseline equivalence of groups, and adequate statistical analyses. Disagreements were resolved by consensus and, if necessary, by the opinion of a third reviewer (M.G.B.).

Statistical analysis

Descriptive statistics were used to summarize the extracted data. Continuous data were presented as mean with standard deviation or as median with (interquartile) range accordingly. Binary or categorical data were presented as frequencies (%). Based on the quality of the data a formal meta-analysis was not performed but rather medians with interquartile range are presented for percentages whenever possible.

Results

After screening 1898 studies, 19 studies with 1338 patients fulfilled the eligibility criteria (Fig. 1), with 177 (13%) patients after MIS LPJ and 1161 (87%) after open LPJ^{25–43} (Table 1). The included studies originated from seven countries (USA, UK, India, the Netherlands, South Korea, France, and Japan). All studies were retrospective in nature, as no prospective observational studies and RCTs were identified. Two studies were comparative studies, whereas 17 were non-comparative. The smallest study was published by Khaled and Ammori²⁹




Fig. 1 PRISMA 2020 flow diagram for included studies

(5 laparoscopic LPJ) and the largest was reported by Napolitano et al.³⁹ (524 open Puestow LPJ). Outcome of MIS LPJ was reported in eight studies, outcome of open LPJ in nine studies, and two studies^{31,32} compared both approaches. Extracted data are displayed in [Tables 1–3](#) and [Tables S1–2](#).

All studies included patients with painful CP ([Table S2](#)). The most common aetiologies for CP were alcohol, tropical, and idiopathic. Postoperative outcome was reported by all studies whereas long-term outcomes such as pain relief, weight gain and new onset of exocrine or endocrine insufficiency were reported by 15, 5 and 7 studies respectively. Pain relief after surgery was evaluated using different methods (for example visual analogue scale, Izbicki pain score, complete or partial pain relief, etc.) and timing of follow-up (1, 2, 3, and 5 years) among studies. The studies by Kirks et al.³¹ and Kalady et al.²⁸ were the only two reporting on costs.

Short-term outcomes in MIS LPJ

Short-term outcomes in MIS LPJ are shown in [Tables 2, 3](#). Operative time per study ranged from 171 to 300 min and the estimated blood loss (EBL) ranged from 42 to 184 ml. Postoperative morbidity rate ranged from 0% to 57% (median and i.q.r. could not be calculated).

Length of hospital stay was from 5 to 7 days, readmission rate from 14% to 38% (median and i.q.r. could not be calculated) and mortality rate was 0% overall (median: 0, i.q.r.: 0).

Short-term outcomes in open LPJ

Short-term outcomes in open LPJ are shown in [Tables 2, 3](#). Operative time varied from 95 to 327 min and EBL from 120 to 600 ml. Postoperative morbidity rate ranged between 4% and 68% (median: 25, i.q.r.: 23). Length of hospital stay ranged from

Table 1 Study characteristics on minimally invasive and open lateral pancreaticojejunostomy for symptomatic chronic pancreatitis

Years	Authors	Country	Study design	MIS versus open	LPJ	Type of approach	MINORS score
Minimally invasive LPJ					(n = 132)		
2004	Tantia et al. ³⁷	India	Retrospective	No	17	Laparoscopic	5/16
2006	Palanivelu et al. ³⁴	India	Retrospective	No	12	Laparoscopic	4/16
2014	Khaled and Ammori ²⁹	UK	Retrospective	No	5	Laparoscopic	8/16
2013	Sahoo and Kumar ³⁵	India	Retrospective	No	12	Laparoscopic	5/16
2016	Kim and Hong ³⁰	Korea	Retrospective	No	11	Laparoscopic	5/16
2018	Hamad et al. ²⁷	USA	Retrospective	No	8	Robotic	10/16
2019	Bhandarwar et al. ²⁵	India	Retrospective	No	28	Laparoscopic	9/16
2019	Senthilnathan et al. ³⁶	India	Retrospective	No	39	Laparoscopic	9/16
Open LPJ					(n = 1104)		
2000	Sielezneff et al. ⁴¹	France	Retrospective	No	57	Open	10/16
2000	Sohn et al. ⁴²	USA	Retrospective	No	52	Open	9/24
2001	Nealon and Matin ⁴⁰	USA	Retrospective	No	124	Open	8/24
2001	Kalady et al. ²⁸	USA	Retrospective	No	60	Open	7/16
2002	Boerma et al. ²⁶	The Netherlands	Retrospective	No	50	Open	8/24
2003	Nealon and Walser ³³	USA	Retrospective	No	103	Open	8/24
2014	Sudo et al. ⁴³	Japan	Retrospective	No	64	Open	7/16
2020	Napolitano et al. ³⁹	USA	Retrospective	No	524	Open	7/16
2022	Kempeneers et al. ³⁸	The Netherlands	Retrospective	No	70	Open	18/24
Minimally invasive versus open LPJ					(n = 102)		
2017	Kirks et al. ³¹	USA	Retrospective	Yes	26	Robotic versus open	14/24
2022	Nag et al. ³²	India	Retrospective	Yes	76	Laparoscopic versus open	18/24

MIS, minimally invasive surgery; LPJ, lateral pancreaticojejunostomy; MINORS, Methodological Index for Non-Randomized Studies.

Table 2 Intraoperative outcomes of lateral pancreaticojejunostomy

Study	Operative time*		EBL†	
	MIS	Open	MIS	Open
Sielezneff et al. ⁴¹	–	–	–	–
Sohn et al. ⁴²	–	312	–	600
Nealon and Matin ⁴⁰	–	–	–	–
Kalady et al. ²⁸	–	–	–	–
Boerma et al. ²⁶	–	S: 170 NS: 165 CJ: 147 CA: 95	–	–
Nealon and Walser ³³	–	–	–	–
Tantia et al. ³⁷	LPJ: 277	–	–	–
Palanivelu et al. ³⁴	179	–	–	–
Khaled and Ammori ²⁹	277	–	150	–
Sahoo and Kumar ³⁵	262	–	–	–
Sudo et al. ⁴³	227	227	–	298
Kirks et al. ³¹	268	236	76	250
Kim and Hong ³⁰	201	–	42	–
Hamad et al. ²⁷	210	–	45	–
Bhandarwar et al. ²⁵	Staple: 171 Sutured: 213	Conversion: 185	–	–
Senthilnathan et al. ³⁶	221	–	184	–
Napolitano et al. ³⁹	–	241	–	–
Nag et al. ³²	300	210	100	120
Kempeneers et al. ³⁸	327	–	300	–

*Mean/median (min); †mean/median (ml); MIS, minimally invasive surgery; S, stent; NS, no stent; CJ, cystojejunostomy; CA, cyst aspiration.

6 to 16 days. Readmission rate ranged from 3% to 33% (median: 16, i.q.r.: 16.5) and in-hospital mortality rate varied from 0% to 3% (median: 0, i.q.r.: 1).

Long-term outcomes

Percentages of the longest follow-up period were used for the calculation of medians and i.q.r. of pain relief and endocrine and exocrine insufficiencies. The rate of pain relief ranged from 62% to 91% after open LPJ (median: 78.5, i.q.r.: 23) and from 71% to 100% (median: 82.5, i.q.r.: 12.5) after MIS LPJ respectively. New-onset endocrine insufficiency was reported in 21% to 22% (median and i.q.r. could not be calculated) in MIS LPJ and for 19% to 26% of patients after open LPJ. New-onset exocrine insufficiency was

shown in 11% to 27% in MIS LPJ versus 8% to 26% after open LPJ (median and i.q.r. could not be calculated). Finally, the rate of weight gain ranged from 60% to 100% (median: 97, i.q.r. 23) after MIS LPJ (Table S1). Weight gain following open LPJ was not reported in the included studies.

Costs

Two studies^{28,31} reported on hospital costs (Table S1). The first study reported a mean hospital cost for open LPJ of \$13 530 whereas the latter compared median total cost between robotic and open LPJ (robotic LPJ \$23 286 versus open LPJ \$27 186)³¹. Additionally, a subanalysis revealed differences in costs for

Table 3 Postoperative outcomes of lateral pancreaticojejunostomy

Study	Morbidity rate†		Length of stay‡		Readmission rate (%)		Mortality rate (%)	
	MIS	Open	MIS	Open	MIS	Open	MIS	Open
Sieleznoff et al. ⁴¹	–	30	–	14	–	–	–	0
Sohn et al. ⁴²	–	31	–	12	–	–	–	2
Nealon and Matin ⁴⁰	–	4	–	–	–	–	–	0
Kalady et al. ²⁸	–	25	–	12	–	3	–	0
Boerma et al. ²⁶	–	S: 19	–	10	–	S: 26	–	0
Nealon and Walser ³³	–	NS: 8	–	CJ 9	–	NS: 33	–	0
		CJ: 16		CA: 9		–		
		CA: 11		–		–		
Tantia et al. ³⁷	12	–	5	–	–	–	0	–
Palanivelu et al. ³⁴	8	–	5	–	–	–	0	–
Khaled and Ammoni ²⁹	20	–	5	–	20	–	0	–
Sahoo and Kumar ³⁵	0	–	8	–	–	–	0	–
Sudo et al. ⁴³	–	33	–	16	–	16	–	0
Kirks et al. ³¹	57	68	5	7	14	16	0	0
Kim and Hong ³⁰	9	–	6	–	–	–	0	–
Hamad et al. ²⁷	25	–	7	–	38	–	0	–
Bhandarwar et al. ²⁵	29	–	6	Conversion: 8.5	–	–	0	–
Senthilnathan et al. ³⁶	POPF: 8 PPH: 3	–	6	–	–	–	0	–
Napolitano et al. ³⁹	–	35	–	10	–	10	–	1
Nag et al. ³²	8	10	7	6	–	–	0	0
Kempeneers et al. ³⁸	–	9	–	9	–	–	–	30-days: 1 90-day: 3

†Mean/median (ml); ‡mean/median (days); MIS, minimally invasive surgery; EBL, estimated blood loss; S, stent; NS, no stent; CJ, cystojejunostomy; CA, cyst aspiration; POPF, postoperative pancreatic fistula; PPH, postpancreatectomy haemorrhage.

operative supplies (robotic LPJ \$4063 versus open LPJ \$1854) and postoperative medication (robotic LPJ \$220 versus open LPJ \$568)³¹.

Methodological quality assessment

Individual MINORS scores in MIS LPJ and open LPJ were relatively low, ranging from 4 to 10 and 7 to 18 respectively. Studies comparing MIS LPJ and open LPJ had higher scores of 14 and 18. RoB2²⁴ was not used as no randomized studies were included.

Discussion

This systematic review on MIS versus open LPJ in patients with symptomatic CP suggests that, based on very low-quality evidence, MIS LPJ may hold some benefits in short-term morbidity rate and length of hospital stay. Long-term outcomes (for example pain relief, exocrine and endocrine insufficiency) appear to be similar between MIS and open LPJ. It must be emphasized that high-quality studies are needed, especially a phase 2 RCT which should also take postoperative pain control and costs into account.

As this is the first systematic review on this topic, these findings could not be compared to previous systematic reviews. Systematic reviews are available for MIS versus open pancreatoduodenectomy, confirming a shorter length of hospital stay with the MIS approach^{44–46}. Compared to pancreatoduodenectomy, LPJ is associated with lower rates of major morbidity and mortality rates, which could be explained by the lower incidence of postoperative pancreatic fistula¹⁶ following LPJ relative to the hard pancreatic texture and atrophic parenchyma resulting from CP. Additionally, the main goal of surgery in CP is to increase QoL by reducing pain.

Because MIS is, in general, associated with less postoperative pain compared to open surgery, it is of paramount importance in the treatment of CP based on the challenges with pain therapy in those patients.

As a result of technological improvements and increasing experience with MIS, surgeons started performing MIS LPJ in the

early 2000s. Some innovations have been reported since. In 2018, Bhandarwar et al.²⁵ reported the feasibility of using staples for performing LPJ, thereby reducing operative time without worsening postoperative outcomes. Key to this technique is the presence of a wide main pancreatic duct to properly place the surgical stapler device. In 2017 the first series from the USA reported on robot-assisted MIS LPJ. Since then, two further studies reported on this approach (at least five patients). The robotic platform may be particularly useful given the extensive amount of suturing that is required during LPJ.

Only two studies directly compared outcome after MIS and open LPJ. These studies, from the USA³¹ and India³², reported largely similar postoperative outcomes after MIS and open LPJ. In a single-centre retrospective non-matched series of 26 patients, Kirks et al. reported lower EBL (75 ml versus 250 ml, $P=0.043$) and length of stay (5 versus 7, $P=0.009$) with robotic MIS LPJ compared to open, despite a 32 min increase in operative time (268 min versus 236 min, $P=0.412$)³¹. Additionally, Kirks et al. reported reduced total cost with the robotic approach (Spearman's rank correlation, robot, $r: -0.775$, $r^2: 0.600$; open, $r: -0.573$, $r^2: 0.328$). In a single-centre matched series of 76 patients, Nag et al. reported lower intraoperative blood loss (100 ml versus 120 ml, $P=0.009$) with laparoscopic MIS LPJ as compared to open LPJ despite a 90 min increased operative time (300 min versus 210 min, $P=0.000$). Notably, this study also reported a high conversion rate of 29%³². These findings should be assessed by further studies. The Miami guidelines for minimally invasive pancreatic surgery provide no guidance on MIS LPJ but advise a minimum volume of 20 MIS pancreatoduodenectomy procedures per centre⁴⁷. Given the lower incidence of LPJ, these numbers are difficult to gather; hence, MIS LPJ might be reserved for high-volume centres, which have experience in both minimally invasive pancreatic surgery and minimally invasive pancreatic surgery specifically for CP.

A crucial point highlighted by the present systematic review was the lack of uniform reporting regarding the assessment of pain relief and the complete absence of reporting on

postoperative (in hospital) pain control. The latter is of obvious importance as this would be an important mechanism for a benefit of MIS LPJ translating into faster time to functional recovery and shorter hospital stay. Regarding long-term pain relief, several definitions were used, making it impossible to perform a proper comparison. Standardized pain assessment with universal pain relief scores should be mandatory to properly understand the potential benefit of surgery in CP patients. In 2020, the Dutch Pancreatitis Study Group published the ESCAPE trial⁸, which reported a superiority of surgery as first intervention for CP compared to endoscopic treatment. Early surgery compared with an endoscopy-first approach resulted in lower pain scores over the course of 18 months. Three years later, Kempeneers *et al.*⁴⁸ further reported results of the ESCAPE trial with early surgery being superior compared to the endoscopy-first approach from a cost-effectiveness point of view. Additionally, performing endoscopy prior to surgery decreases the benefit of surgery⁴⁹. Based on these and other studies, the role of surgery as reference standard for chronic pancreatitis is now widely accepted. However, proper standardization of intraoperative management, postoperative management and follow-up is still lacking.

Several limitations should be considered when interpreting the results of this systematic review. There was considerable heterogeneity between the included studies in terms of outcome reporting, thereby making results difficult to interpret. Future studies should report complications in accordance with current guidelines (for example ISGPS definitions^{16–21}) and define outcomes properly in the study methods. The study size varies remarkably between studies on MIS (median 12 patients) and open (median 64 patients) LPJ. This was driven by the inclusion criteria of minimum 5 MIS LPJ or 50 open LPJ, highlighting a significant bias related to the quality of data because the open group holds more data from high-volume centres and thus likely reporting better quality. Several included studies report follow-up times that are not adequate for patients with CP because pain relief is not always expected directly following surgery but is expected in the years after surgery, so future studies including a minimum of 12 months of follow-up are necessary. Although 177 patients after MIS LPJ were included in this review, only 15 patients underwent robot-assisted MIS LPJ. Clearly more data on this subgroup are needed. Literature comprised retrospective studies with a relatively low number of patients with CP and relatively low methodological quality resulting in a systematic review of low-quality evidence only. According to the IDEAL recommendations⁵⁰, pragmatic phase 2 RCTs are now warranted to provide stronger evidence on the role of MIS LPJ in eligible patients with symptomatic CP. Most MIS LPJ studies (75%) were published after 2014 whereas the majority of open LPJ studies (67%) were published before 2004. This highlighted a strong temporal bias, as surgical tools and postoperative management have improved in the last decade.

In conclusion, this systematic review of low-quality evidence suggested that MIS LPJ can be performed safely in selected patients. The next step would be to perform a phase 2 RCT with standardized outcome measures focusing on potential short-term benefits for patients.

Author contributions

Roberto M. Montorsi (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing—original draft,

Writing—review & editing), Michiel F.G. Francken (Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing—original draft, Writing—review & editing), Marja A. Boermeester (Validation, Visualization, Writing—review & editing), Olivier R. Busch (Validation, Visualization, Writing—review & editing), Freek Daams (Validation, Visualization, Writing—review & editing), Thilo Hackert (Validation, Visualization, Writing—review & editing), Roel Haen (Validation, Visualization, Writing—review & editing), Markus W. Hollmann (Validation, Visualization, Writing—review & editing), Hjalmar C. van Santvoort (Validation, Visualization, Writing—review & editing), and Marc G. Besselink (Conceptualization, Supervision, Validation, Visualization, Writing—review & editing)

Funding

The authors have no funding to declare.

Acknowledgements

R.M.M. and M.F.G.F. are shared first authorship.

Disclosure

The authors declare no conflict of interest.

Supplementary material

[Supplementary material](#) is available at *BJS Open* online.

Data availability

No new data were generated or analyzed in support of this research.

References

- Whitcomb DC, Frulloni L, Garg P, Greer JB, Schneider A, Yadav D *et al.* Chronic pancreatitis: an international draft consensus proposal for a new mechanistic definition. *Pancreatology* 2016; **16**:218–224
- Hoffmeister A, Mayerle J, Beglinger C, Büchler MW, Bufler P, Dathe K *et al.* English language version of the S3-consensus guidelines on chronic pancreatitis: definition, aetiology, diagnostic examinations, medical, endoscopic and surgical management of chronic pancreatitis. *Z Gastroenterol* 2015; **53**: 1447–1495
- Drewes AM, Bouwense SAW, Campbell CM, Ceyhan GO, Delhaye M, Demir IE *et al.* Guidelines for the understanding and management of pain in chronic pancreatitis. *Pancreatology* 2017; **17**:720–731
- Poulsen JL, Olesen SS, Malver LP, Frøkjær JB, Drewes AM. Pain and chronic pancreatitis: a complex interplay of multiple mechanisms. *World J Gastroenterol* 2013; **19**:7282–7291
- Machicado JD, Chari ST, Timmons L, Tang G, Yadav D. A population-based evaluation of the natural history of chronic pancreatitis. *Pancreatology* 2018; **18**:39–45
- Cahen DL, Gouma DJ, Nio Y, Rauws EAJ, Boermeester MA, Busch OR *et al.* Endoscopic versus surgical drainage of the pancreatic duct in chronic pancreatitis. *N Engl J Med* 2007; **356**:676–684

7. Díte P, Ruzicka M, Zboril V, Novotný I. A prospective, randomised trial comparing endoscopic and surgical therapy for chronic pancreatitis. *Endoscopy* 2003;**35**:553–558
8. Issa Y, Kempeneers MA, Bruno MJ, Fockens P, Poley JW, Ahmed Ali U et al. Effect of early surgery vs endoscopy-first approach on pain in patients with chronic pancreatitis: the ESCAPE randomised clinical trial. *JAMA* 2020;**323**:237–247
9. Löhr JM, Dominguez-Munoz E, Rosendahl J, Besselink M, Mayerle J, Lerch MM et al. United European gastroenterology evidence-based guidelines for the diagnosis and therapy of chronic pancreatitis (HaPanEU). *United European Gastroenterol J* 2017;**5**:153–199
10. Gardner TB, Adler DG, Forsmark CE, Sauer BG, Taylor JR, Whitcomb DC. ACG clinical guideline: chronic pancreatitis. *Am J Gastroenterol* 2020;**115**:322–339
11. Shimizu K, Ito T, Irisawa A, Ohtsuka T, Ohara H, Kanno A et al. Evidence-based clinical practice guidelines for chronic pancreatitis 2021. *J Gastroenterol* 2022;**57**:709–724
12. Kempeneers MA, Issa Y, Ali UA, Baron RD, Besselink MG, Büchler M et al. International consensus guidelines for surgery and the timing of intervention in chronic pancreatitis. *Pancreatol* 2020;**20**:149–157
13. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021;**88**:105906
14. Booth A, Clarke M, Dooley G, Ghera D, Moher D, Petticrew M et al. The nuts and bolts of PROSPERO: an international prospective register of systematic reviews. *Syst Rev* 2012;**1**:2
15. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 2016;**5**:210
16. Bassi C, Marchegiani G, Dervenis C, Sarr M, Abu Hilal M, Adham M et al. The 2016 update of the international study group (ISGPS) definition and grading of postoperative pancreatic fistula: 11 years after. *Surgery* 2017;**161**:584–591
17. Wente MN, Bassi C, Dervenis C, Fingerhut A, Gouma DJ, Izbicki JR et al. Delayed gastric emptying (DGE) after pancreatic surgery: a suggested definition by the International Study Group of Pancreatic Surgery (ISGPS). *Surgery* 2007;**142**:761–768
18. Wente MN, Veit JA, Bassi C, Dervenis C, Fingerhut A, Gouma DJ et al. Postpancreatectomy hemorrhage (PPH): an International Study Group of Pancreatic Surgery (ISGPS) definition. *Surgery* 2007;**142**:20–25
19. Koch M, Garden OJ, Padbury R, Rahbari NN, Adam R, Capussotti L et al. Bile leakage after hepatobiliary and pancreatic surgery: a definition and grading of severity by the International Study Group of Liver Surgery. *Surgery* 2011;**149**:680–688
20. Besselink MG, van Rijssen LB, Bassi C, Dervenis C, Montorsi M, Adham M et al. Definition and classification of chyle leak after pancreatic operation: a consensus statement by the International Study Group on Pancreatic Surgery. *Surgery* 2017;**161**:365–372
21. Marchegiani G, Barreto SG, Bannone E, Sarr M, Vollmer CM, Connor S et al. Postpancreatectomy acute pancreatitis (PPAP): definition and grading from the International Study Group for Pancreatic Surgery (ISGPS). *Ann Surg* 2022;**275**:663–672
22. Clavien PA, Barkun J, de Oliveira ML, Vauthey JN, Dindo D, Schulick RD et al. The Clavien–Dindo classification of surgical complications: five-year experience. *Ann Surg* 2009;**250**:187–196
23. Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomised studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003;**73**:712–716
24. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al. RoB2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;**366**:14898
25. Bhandarwar A, Arora E, Gajbhiye R, Gandhi S, Patel C, Wagh A et al. Laparoscopic lateral pancreaticojejunostomy: an evolution to endostapled technique. *Surg Endosc* 2019;**33**:1749–1756
26. Boerma D, van Gulik TM, Rauws EA, Obertop H, Gouma DJ. Outcome of pancreaticojejunostomy after previous endoscopic stenting in patients with chronic pancreatitis. *Eur J Surg* 2002;**168**:223–228
27. Hamad A, Zenati MS, Nguyen TK, Hogg ME, Zeh HJ, Zureikat AH. Safety and feasibility of the robotic platform in the management of surgical sequelae of chronic pancreatitis. *Surg Endosc* 2018;**32**:1056–1065
28. Kalady MF, Broome AH, Meyers WC, Pappas TN. Immediate and long-term outcomes after lateral pancreaticojejunostomy for chronic pancreatitis. *Am Surg* 2001;**67**:478–483
29. Khaled Y, Ammori B. Laparoscopic lateral pancreaticojejunostomy and laparoscopic Berne modification of Beger procedure for the treatment of chronic pancreatitis: the first UK experience. *Surg Laparosc Endosc Percutan Tech* 2014;**24**:e178–e182
30. Kim EY, Hong TH. Laparoscopic longitudinal pancreaticojejunostomy using barbed sutures: an efficient and secure solution for pancreatic duct obstructions in patients with chronic pancreatitis. *J Gastrointest Surg* 2016;**20**:861–866
31. Kirks RC, Lorimer PD, Fruscione M, Cochran A, Baker EH, Iannitti DA et al. Robotic longitudinal pancreaticojejunostomy for chronic pancreatitis: comparison of clinical outcomes and cost to the open approach. *Int J Med Robot* 2017;**13**:e1832
32. Nag H, Nekarakanti P, Arvinda P, Sharma A. Laparoscopic versus open surgical management of patients with chronic pancreatitis: a matched case–control study. *J Minim Access Surg* 2022;**18**:191–196
33. Nealon WH, Walser E. Duct drainage alone is sufficient in the operative management of pancreatic pseudocyst in patients with chronic pancreatitis. *Ann Surg* 2003;**237**:614–620, discussion 620–622
34. Palanivelu C, Shetty R, Jani K, Rajan PS, Sendhil Kumar K, Parthasarathi R et al. Laparoscopic lateral pancreaticojejunostomy: a new remedy for an old ailment. *Surg Endosc* 2006;**20**:458–461
35. Sahoo M, Kumar A. Laparoscopic longitudinal pancreaticojejunostomy using cystoscope and ERCP basket for retrieval of leftover pancreatic duct stones. *Surg Endosc* 2013;**27**:S288
36. Senthilnathan P, Subrahmaneswara Babu N, Vikram A, Sabnis SC, Srivatsan Gurumurthy S, Anand Vijai N et al. Laparoscopic longitudinal pancreaticojejunostomy and modified Frey's operation for chronic calcific pancreatitis. *BJS Open* 2019;**3**:666–671
37. Tania O, Jindal MK, Khanna S, Sen B. Laparoscopic lateral pancreaticojejunostomy: our experience of 17 cases. *Surg Endosc* 2004;**18**:1054–1057
38. Kempeneers MA, van Hemert AKE, van der Hoek M, Issa Y, van Hooft JE, Nio CY et al. Short- and long-term outcomes of selective use of Frey or extended lateral pancreaticojejunostomy in chronic pancreatitis. *Br J Surg* 2022;**109**:363–371
39. Napolitano M, Brody F, Lee KB, Rosenfeld E, Chen S, Murillo-Berlitz AE et al. 30-day outcomes and predictors of complications after Puestow procedure. *Am J Surg* 2020;**220**:372–375
40. Nealon WH, Matin S. Analysis of surgical success in preventing recurrent acute exacerbations in chronic pancreatitis. *Ann Surg* 2001;**233**:793–800
41. Sielezneff I, Malouf A, Salle E, Brunet C, Thirion X, Sastre B. Long term results of lateral pancreaticojejunostomy for chronic alcoholic pancreatitis. *Eur J Surg* 2000;**166**:58–64

42. Sohn TA, Campbell KA, Pitt HA, Sauter PK, Coleman JA, Lillemo KD et al. Quality of life and long-term survival after surgery for chronic pancreatitis. *J Gastrointest Surg* 2000;**4**:355–364, discussion 364–365
43. Sudo T, Murakami Y, Uemura K, Hashimoto Y, Kondo N, Nakagawa N et al. Short- and long-term results of lateral pancreaticojejunostomy for chronic pancreatitis: a retrospective Japanese single-center study. *J Hepatobiliary Pancreat Sci* 2014;**21**:426–432
44. de Rooij T, Lu MZ, Steen MW, Gerhards MF, Dijkgraaf MG, Busch OR et al. Minimally invasive versus open pancreatoduodenectomy: systematic review and meta-analysis of comparative cohort and registry studies. *Ann Surg* 2016;**264**:257–267
45. Correa-Gallego C, Dinkelspiel HE, Sulimanoff I, Fisher S, Viñuela EF, Kingham TP et al. Minimally-invasive vs open pancreaticoduodenectomy: systematic review and meta-analysis. *J Am Coll Surg* 2014;**218**:129–139
46. van der Heijde N, Balduzzi A, Alseidi A, Dokmak S, Polanco PM, Sandford D et al. The role of older age and obesity in minimally invasive and open pancreatic surgery: a systematic review and meta-analysis. *Pancreatology* 2020;**20**:1234–1242
47. Asbun HJ, Moekotte AL, Vissers FL, Kunzler F, Cipriani F, Alseidi A et al. The Miami International Evidence-based Guidelines on Minimally Invasive Pancreas Resection. *Ann Surg* 2020;**271**:1–14
48. Kempeneers MA, Issa Y, Bruno MJ, van Santvoort HC, Besselink MG, Boermeester MA et al. Cost-effectiveness of early surgery versus endoscopy-first approach for painful chronic pancreatitis in the ESCAPE trial. *Ann Surg* 2023;**277**:e878–e884
49. Ahmed Ali U, Nieuwenhuijs VB, van Eijck CH, Gooszen HG, van Dam RM, Busch OR et al. Clinical outcome in relation to timing of surgery in chronic pancreatitis: a nomogram to predict pain relief. *Arch Surg* 2012;**147**:925–932
50. McCulloch P, Altman DG, Campbell WB, Flum DR, Glasziou P, Marshall JC et al. No surgical innovation without evaluation: the IDEAL recommendations. *Lancet* 2009;**374**:1105–1112