

Laryngeal and Pharyngeal Squamous Cell Carcinoma After Antireflux Surgery in the 5 Nordic Countries

Manar Yanes, MD,* Giola Santoni, PhD,* John Maret-Ouda, MD, PhD,*†
 Eivind Ness-Jensen, MD, PhD,*‡§ Martti Färkkilä, MD, PhD,||
 Elsebeth Lyngge, MD, PhD,¶ Eero Pukkala, PhD,#** Pål Romundstad, MSc, PhD,‡
 Laufey Tryggvadóttir, MSc,††‡‡ My von Euler-Chelpin, PhD,§§
 and Jesper Lagergren, MD, PhD*|||✉

Objective: The aim of this study was to clarify whether antireflux surgery prevents laryngeal and pharyngeal squamous cell carcinoma.

Summary Background Data: Gastroesophageal reflux disease (GERD) seems to increase the risk of laryngeal and pharyngeal squamous cell carcinoma.

Methods: All-Nordic (Denmark, Finland, Iceland, Norway, and Sweden) population-based cohort study of adults with documented GERD in 1980 to 2014. First, cancer risk after antireflux surgery was compared to the expected risk in the corresponding background population by calculating standardized incidence ratios (SIRs) with 95% confidence intervals (CIs). Second, cancer risk among antireflux surgery patients was compared to nonoperated GERD patients using multivariable Cox regression, providing hazard ratios (HR) with 95% CIs, adjusted for sex, age, calendar period, and diagnoses related to tobacco smoking, obesity, and alcohol overconsumption.

Results: Among 814,230 GERD patients, 47,016 (5.8%) underwent antireflux surgery. The overall SIRs and HRs of the combined outcome

laryngeal or pharyngeal squamous cell carcinoma (n=39) were decreased after antireflux surgery [SIR=0.62 (95% CI 0.44–0.85) and HR=0.55 (95% CI 0.38–0.80)]. The point estimates were further decreased >10 years after antireflux surgery [SIR=0.48 (95% CI 0.26–0.80) and HR=0.47 (95% CI 0.26–0.85)]. The risk estimates of laryngeal squamous cell carcinoma were particularly decreased >10 years after antireflux surgery [SIR=0.28 (95% CI 0.08–0.72) and HR=0.23 (95% CI 0.08–0.69)], whereas no such decrease over time after surgery was found for pharyngeal squamous cell carcinoma. Analyses of patients with severe GERD (reflux esophagitis or Barrett esophagus) showed similar results.

Conclusion: Antireflux surgery may decrease the risk of laryngeal squamous cell carcinoma and possibly also of pharyngeal squamous cell carcinoma.

Keywords: antireflux medication, fundoplication, gastroesophageal reflux disease, larynx, neoplasm, Nissen, pharynx, proton pump inhibitor

(*Ann Surg* 2022;276:e79–e85)

From the *Upper Gastrointestinal Surgery, Department of Molecular medicine and Surgery, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden; †Centre for Clinical Research Sörmland, Uppsala University, Eskilstuna, Sweden; ‡Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim/Levanger, Norway; §Medical Department, Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway; ||Clinic of Gastroenterology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland; ¶Nykøbing Falster Hospital, University of Copenhagen, Denmark; #Finnish Cancer Registry, Institute for Statistical and Epidemiological Cancer Research, Helsinki, Finland; **Faculty of Social Sciences, Tampere University, Tampere, Finland; ††Icelandic Cancer Registry, Icelandic Cancer Society, Reykjavik, Iceland; ‡‡Faculty of Medicine, University of Iceland, Reykjavik, Iceland; §§Department of Public Health, University of Copenhagen, Copenhagen, Denmark; and |||School of Cancer and Pharmaceutical Sciences, King's College London, United Kingdom.

✉Jesper.Lagergren@ki.se.

Author's contribution: Study concept and design were done by all authors; acquisition of data was done by all authors; analysis and interpretation of data were done by M.Y., G.S., and J.L.; drafting of the manuscript was done by M.Y. and J.L.; critical revision of the manuscript for important intellectual content was done by all authors; statistical analysis was done by M.Y. and G.S.; funding was obtained by J.L.; study supervision was done by J.L.

Funding: This work was supported by the Nordic Cancer Union (grant number 186058), Swedish Cancer Society (grant number 180684), and Swedish Research Council (grant number 340–2013–5478). Jesper Lagergren was supported by the United European Gastroenterology Research Prize and the Distinguished Professor Award at Karolinska Institutet.

The authors report no conflicts of interest.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.annalsurgery.com).

Copyright © 2020 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 0003-4932/22/27602-0e79

DOI: 10.1097/SLA.0000000000004423

Gastroesophageal reflux disease (GERD), defined by troublesome and long-lasting symptoms of heartburn or regurgitation, affects approximately 20% of adults in Western populations.^{1,2} Long-standing GERD can damage the esophageal mucosa, and cause esophagitis, metaplasia (Barrett esophagus), and esophageal adenocarcinoma.^{3,4} GERD can also lead to more proximal reflux, that is, reflux reaching the larynx and pharynx, where it might cause mucosal injury, inflammation, and tumor development.⁵ Interestingly, the laryngeal and pharyngeal squamous cell epithelium seems to be more susceptible to damage by duodenogastric contents than the esophageal squamous cell epithelium.⁶ Thus, GERD appears to be associated with an increased risk of laryngeal and pharyngeal squamous cell carcinoma.^{5,7–12} GERD-related carcinogenesis in the esophagus depends on the development of Barrett esophagus as an intermediate step before adenocarcinoma, and is not associated with esophageal squamous cell carcinoma.¹³ Laryngeal and pharyngeal squamous cell carcinomas are otherwise mainly associated with tobacco smoking and high consumption of alcohol.¹⁴ New preventive strategies could arise if antireflux treatment prevents GERD-associated cancers. Antireflux surgery is a valuable tool for research purposes in this respect because it anatomically and physiologically hinders or reduces reflux starting from a specific date. Yet, no study has evaluated whether the risk of laryngeal or pharyngeal squamous cell carcinoma decreases after antireflux surgery, which was the aim of the present multi-country and population-based cohort study.

METHODS

Study Design

This population-based cohort study in the 5 Nordic countries (Denmark, Finland, Iceland, Norway, and Sweden) investigated associations between antireflux surgery in patients with GERD (exposure, defined in Supplementary Table 1, <http://links.lww.com/SLA/C560>) and the risk of laryngeal and pharyngeal squamous cell carcinoma (outcomes, defined in Supplementary Table 2, <http://links.lww.com/SLA/C560>) during the study period 1980 through 2014. We included only squamous cell carcinomas because other histological types are rare and have different etiologies. The risk of laryngeal and pharyngeal squamous cell carcinoma among patients who have undergone antireflux surgery was compared with 2 groups: the background Nordic population and GERD patients who did not undergo antireflux surgery.

Nordic Antireflux Surgery Cohort

The cohort, entitled the Nordic Antireflux Surgery Cohort (NordASCO), was created by merging data from well-established nationwide health data registries with a long history in the Nordic countries. This cohort has been described in detail elsewhere.¹⁵ In brief, all cohort participants had a diagnosis of GERD documented in a patient registry in any of the Nordic countries. A subcohort only included patients with severe GERD, that is, those with and objectively defined reflux esophagitis or Barrett esophagus. The overall study period was from January 1, 1980 to December 31, 2014, but the start and end years differed between the countries. All participants were aged ≥ 18 and ≤ 95 years at the time of the first GERD diagnosis. We excluded individuals with laryngeal or pharyngeal squamous cell carcinoma before the diagnosis of GERD. The similarity in structure of the Nordic countries' health data registries enabled merging of the data into 1 large cohort. The personal identity numbers of each resident in the Nordic countries allowed correct linkages of individuals' data between registries.^{15,16} Ethical and data permissions were retrieved from the relevant authorities within each country.¹⁵

Data Collection

The data for the patients in the GERD cohort came from the nationwide patient registries, cancer registries, and cause of death registries in the Nordic countries. The data for the background population came from the national total population registries and cancer registries. To assess the use of medications in nonoperated patients with GERD, we retrieved data from the Swedish Prescribed Drug Registry. These 5 types of registries are presented below.

The patient registries were used to identify all cohort participants with a documented diagnosis of GERD, including those who underwent an antireflux surgery procedure of fundoplication for GERD, by providing diagnosis codes and surgical procedure codes from in-hospital and specialized outpatient care in each Nordic country. The specific codes used for defining GERD, severe GERD (reflux esophagitis or Barrett's esophagus), and antireflux surgery in the patient registries are presented in Supplementary Table 1, <http://links.lww.com/SLA/C560>. Additional data collected from these registries for the present study were sex, date of birth, and diagnoses associated with tobacco smoking (chronic obstructive pulmonary disease), obesity (obesity diagnosis and diabetes type 2), and alcohol overconsumption (alcohol-associated vitamin B deficiency, liver disease, and pancreatic disease). Complete nationwide coverage

of the patient registries was reached in 1967 (Finland), 1978 (Denmark), 1987 (Sweden), 1997 (Norway), and 1999 (Iceland). The data recorded in these registries have high positive predictive values for most diagnoses and surgical procedures.^{17–19}

The cancer registries were used to identify participants who developed laryngeal or pharyngeal squamous cell carcinoma (Supplementary Table 2, <http://links.lww.com/SLA/C560>). The collected data included date of cancer diagnosis, histological type and location of the tumor, and age at cancer diagnosis. All Nordic cancer registries have been nationwide since their initiation in 1943 (Denmark), 1953 (Norway), 1953 (Finland), 1955 (Iceland), and 1958 (Sweden). Numerous validation studies have shown high completeness ($\geq 98\%$) and accuracy ($\geq 94\%$) of the cancer registration.²⁰

The cause of death registries provided death dates with virtually 100% completeness and accuracy.^{16–21} These registries enabled censoring of GERD patients in the cohort who died during follow-up.

The registries of the total population combined with the cancer registries provided data on population count and number of laryngeal and pharyngeal squamous cell carcinoma in the general background populations by age, sex and calendar year in each Nordic country.

The Swedish Prescribed Drug Registry provided data on prescriptions of proton pump inhibitors and histamine-2-receptor antagonists, which were used to validate the use of antireflux medication among nonoperated patients with GERD. The nationwide registration started in July 1, 2005.

Statistical Analysis

Patients in the GERD cohort contributed person-time at risk of laryngeal or pharyngeal squamous cell carcinoma from the date of the first GERD diagnosis or the date of the first antireflux surgery for GERD. The first year of follow-up was excluded to avoid detection bias, that is, earlier detection of any laryngeal or pharyngeal squamous cell carcinoma because of the GERD diagnosis or the antireflux surgery. Person-time at risk in the antireflux surgery group was accumulated from 1 year after the date of surgery until the first occurrence of laryngeal or pharyngeal squamous cell carcinoma, death, or end of study period. Person-time at risk in the nonoperated group with GERD was accumulated from > 1 year after the date of GERD diagnosis until the date of laryngeal or pharyngeal squamous cell carcinoma, death, end of study period, or antireflux surgery. Thus, GERD patients who underwent antireflux surgery during follow-up were censored from the nonoperated GERD group at the date of surgery, and included in the antireflux surgery group instead.

Two statistical approaches were used. First, the cancer risk in the antireflux surgery group (and the nonoperated GERD group) was compared to the corresponding background population by calculating standardized incidence ratios (SIRs) and 95% confidence intervals (CIs). For each group, the observed number of laryngeal or pharyngeal squamous cell carcinoma was divided by the expected number among individuals of the corresponding country, sex (men or women), age (5-year categories), and calendar period (5-year categories). The expected number for each stratum was calculated by multiplying the number of person-years by the respective incidence rate derived from the national cancer registry data. SIRs were computed for the entire follow-up time (> 1 –34 years) as well as for each of the follow-up categories > 1 to 5, > 5 to 10, and > 10 years.

In the second statistical approach, the cancer risk in the antireflux surgery group was compared with that of the nonoperated group with GERD. Multivariable Cox regression was

used to calculate hazard ratios (HR) and 95% CIs, adjusted for sex (men or women), age (continuous), calendar period (1980–1989, 1990–1999, or 2000–2014), country (Denmark, Finland, Iceland, Norway, or Sweden), chronic obstructive pulmonary disease (yes or no), obesity diagnosis or diabetes mellitus type 2 (yes or no), and alcohol-related diagnoses (yes or no). The overall HR assessed the entire period (> 1–34 years), whereas the follow-up categories of HR were > 1 to 5, > 5 to 10, and >10 years.

We also conducted analyses restricted to patients with severe GERD, that is, reflux esophagitis or Barrett esophagus, in which follow-up started from 1 year after the date of the first severe GERD diagnosis or the date of antireflux surgery for severe GERD. Norwegian GERD patients were excluded from these subanalyses because 4-character subcategories of diagnosis codes were not available in the Norwegian patient registry, making it impossible to distinguish severe GERD from any GERD.

In a sensitivity analysis, we evaluated potential bias from selection or confounding by using the above-mentioned

multivariable Cox regression model to assess the risk of colon cancer in the antireflux surgery group compared to the nonoperated group with GERD. Colon cancer was analyzed because it is unrelated to GERD and antireflux surgery and is the most common cancer that occurs frequently in both sexes in the Nordic countries.²²

To validate the use of antireflux medication in the nonoperated group with GERD, we assessed the use of proton pump inhibitors or histamine-2-receptor antagonists in Swedish patients with GERD from 2007 onwards. Nationwide registration of medications started in July 1, 2005, but this validation started from 2007 to exclude ongoing medication and accurately assess new treatment episodes. Among nonoperated cohort participants, 183,699 (92.1%) patients with any GERD and 7143 (97.3%) with severe GERD had dispensed prescriptions of antireflux medication. Among the patients with such prescriptions, 165,773 (90.2%) patients with any GERD and 6530 (91.4%) with severe GERD obtained their first prescription within 3 months before or after first diagnosis of GERD.

TABLE 1. Characteristics of Patients With Gastroesophageal Reflux Disease Having Undergone Antireflux Surgery or Not

Gastroesophageal Reflux Disease		
	No Antireflux Surgery, No. (%)	Antireflux Surgery, No. (%)
Total		
Patients*	780,546 (100)	47,016 (100)
Person-years of follow-up	5,020,529	556,234
Sex		
Male	378,757 (48.5)	26,478 (56.3)
Female	401,789 (51.5)	20,538 (43.7)
Age at inclusion, y		
< 50	256,475 (32.9)	22,095 (47.0)
50–< 65	237,604 (30.4)	18,226 (38.8)
≥65	286,467 (36.7)	6695 (14.2)
Obesity diagnosis	38,889 (5.0)	2267 (5.5)
Diabetes mellitus type 2	78,385 (10.0)	3959 (8.4)
Chronic obstructive pulmonary disease	70,541 (9.0)	3822 (8.1)
Excessive alcohol consumption	50,355 (6.5)	3508 (7.5)
Vitamin B deficiency associated with alcohol	2163 (0.3)	92 (0.2)
Liver and pancreatic disease related to alcohol intake	15,663 (2.0)	718 (1.5)
Laryngeal or pharyngeal squamous cell carcinoma	699 (0.09)[100]	39 (0.08)[100]
Laryngeal squamous cell carcinoma	311 [44.5]	20 [51.3]
Pharyngeal squamous cell carcinoma	388 [55.5]	19 [48.7]
Severe Gastroesophageal Reflux Disease (With Reflux Esophagitis or Barrett Esophagus)		
	No Antireflux Surgery, No. (%)	Antireflux Surgery, No. (%)
Total		
Patients	242,619 (100)	34,766 (100)
Person-years of follow-up	1,996,651	425,331
Sex		
Male	133,827 (55.2)	20,065 (57.7)
Female	108,792 (44.8)	14,701 (42.3)
Age at inclusion, y		
< 50	76,813 (31.7)	16,737 (48.1)
50–< 65	76,918 (31.7)	13,768 (39.6)
≥65	88,888 (36.6)	4261 (12.3)
Obesity diagnosis	14,267 (5.9)	1670 (4.8)
Diabetes mellitus type 2	30,649 (12.6)	3023 (8.7)
Chronic obstructive pulmonary disease	26,906 (11.1)	2798 (8.0)
Excessive alcohol consumption	23,428 (9.7)	2824 (8.1)
Vitamin B deficiency associated with alcohol	1087 (0.4)	69 (0.2)
Liver and pancreatic disease related to alcohol intake	7678 (3.2)	584 (1.7)
Laryngeal or pharyngeal squamous cell carcinoma	321 (0.13)[100]	28 (0.08)[100]
Laryngeal squamous cell carcinoma	148 [46.1]	12 [42.9]
Pharyngeal squamous cell carcinoma	173 [53.9]	16 [57.1]

*Among the nonoperated patients, 13,332 were also included in the operated group after they were censored from the nonoperated group at the date of admission to antireflux surgery.

The data management and statistical analyses followed a predefined study protocol, and were conducted using IBM SPSS Statistics version 24 (IBM Corp, Armonk, NY).

RESULTS

Study Participants

Among all 814,230 cohort participants with GERD, 47,016 (5.8%) underwent antireflux surgery during the study period. The person-years at risk were 556,234 in the antireflux surgery group and 5,020,529 in the nonoperated group with GERD. The subcohort of patients with severe GERD consisted of 269,656 individuals, including 34,766 (12.9%) who underwent antireflux surgery. Table 1 shows some characteristics of the study participants with GERD who underwent antireflux surgery and not. The proportion of men was greater (56.3%) in the antireflux surgery group than in the nonoperated group (48.5%), and the median age at inclusion was lower in the antireflux surgery group [51 years, interquartile range (IQR) 41–59 years] compared to the nonoperated GERD group (58 years, IQR 45–70 years). During follow-up, 39 (0.08%) patients in the antireflux surgery group and 699 (0.09%) in the nonoperated group with GERD developed laryngeal or pharyngeal squamous cell carcinoma. In the subcohort of severe GERD, 28 (0.08%) and 321 (0.13%) patients developed any of these tumors in the antireflux surgery and non-operated groups, respectively.

Operated Patients Compared With the Background Population

After antireflux surgery for GERD, the overall SIRs of the combined outcome laryngeal or pharyngeal squamous cell carcinoma was decreased (SIR 0.62, 95% CI 0.44–0.85), and the point estimate was further decreased after > 10 years of follow-up (SIR 0.48, 95% CI 0.26–0.80). In a separate analysis of laryngeal squamous cell carcinoma, the overall SIR was 0.65 (95% CI 0.52–1.32), and > 10 years after surgery the SIR decreased to 0.28 (95% CI 0.08–0.72) (Table 2). The overall SIR of pharyngeal squamous cell carcinoma was also decreased (SIR 0.60, 95% CI 0.36–0.93), but the risk estimates did not decrease

further with longer follow-up time after antireflux surgery (Table 2). The subanalysis of individuals with severe GERD revealed similar results (Table 3).

Nonoperated Patients Compared With the Background Population

In the nonoperated group with GERD, the risk of laryngeal or pharyngeal squamous cell carcinoma was similar to that of the corresponding background population (overall SIR 1.00, 95% CI 0.93–1.08, n=699), without differences between the 2 tumors or over follow-up (Table 2). The subanalyses of severe GERD showed similar results (Table 3).

Operated Compared With Nonoperated Patients

Compared to nonoperated patients with GERD, those who had undergone antireflux surgery had a decreased overall adjusted HR of the combined outcome laryngeal or pharyngeal squamous cell carcinoma (HR 0.55, 95% CI 0.38–0.80), and the point estimate was further slightly reduced after > 10 years of follow-up (HR 0.47, 95% CI 0.26–0.85) (Table 4). The overall adjusted HR of laryngeal squamous cell carcinoma analyzed separately was decreased after antireflux surgery (HR 0.53, 95% CI 0.31–0.90), and the point estimate was further decreased after > 10 years of follow-up (HR 0.23, 95% CI 0.08–0.69) (Table 4). The overall HR of pharyngeal squamous cell carcinoma was also decreased after antireflux surgery (adjusted HR 0.58, 95% CI 0.35–0.96), but the risk estimates did not decrease with longer follow-up (Table 4). In analyses restricted to patients with severe GERD, the results were mainly similar, but the adjusted point estimates for laryngeal squamous cell carcinoma were further decreased for the overall risk (HR 0.42, 95% CI 0.21–0.83) and for the risk after > 10 years of follow-up (HR 0.19, 95% CI 0.05–0.69) (Table 5).

The sensitivity analysis showed no decreased overall risk of colon cancer in patients GERD who had undergone antireflux surgery compared to those not operated (adjusted HR 1.16, 95% CI 1.04–1.30), and the risk was not decreased in any of the follow-up periods (Supplementary Table 3, <http://links.lww.com/SLA/C560>).

TABLE 2. Risk of Laryngeal and Pharyngeal Squamous Cell Carcinoma Among Patients With Gastroesophageal Reflux Disease Having Undergone Antireflux Surgery or Not, Compared to the Risk in a Corresponding Background Population, Presented as SIR With 95% 95% CI

No Antireflux Surgery								
Follow-Up, y	Total (n)	Person-Years	Laryngeal or Pharyngeal Squamous Cell Carcinoma		Laryngeal Squamous Cell Carcinoma		Pharyngeal Squamous Cell Carcinoma	
			Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)
> 1–34	780,546	5,020,529	699	1.00 (0.93–1.08)	311	0.94 (0.84–1.05)	388	1.06 (0.96–1.17)
> 1–5	780,546	2,410,469	305	1.05 (0.94–1.17)	133	0.95 (0.79–1.12)	172	1.15 (0.98–1.33)
> 5–10	438,408	1,534,799	212	0.99 (0.87–1.14)	99	0.99 (0.80–1.20)	113	1.00 (0.83–1.20)
> 10	212,641	1,075,262	182	0.95 (0.81–1.09)	79	0.88 (0.70–1.10)	103	1.00 (0.82–1.21)
Antireflux Surgery								
Follow-Up, y	Total (n)	Person-Years	Laryngeal or Pharyngeal Squamous Cell Carcinoma		Laryngeal Squamous Cell Carcinoma		Pharyngeal Squamous Cell Carcinoma	
			Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)
> 1–34	47,016	556,234	39	0.62 (0.44–0.85)	20	0.65 (0.40–1.00)	19	0.60 (0.36–0.93)
> 1–5	47,016	176,353	8	0.54 (0.23–1.07)	5	0.67 (0.22–1.57)	3	0.41 (0.08–1.20)
> 5–10	40,649	179,639	17	0.91 (0.53–1.45)	11	1.20 (0.60–2.14)	6	0.63 (0.23–1.37)
> 10	30,159	200,242	14	0.48 (0.26–0.80)	4	0.28 (0.08–0.72)	10	0.67 (0.32–1.23)

TABLE 3. Risk of Laryngeal and Pharyngeal Squamous Cell Carcinoma Among Patients With Severe Gastroesophageal Reflux Disease (With Reflux Esophagitis or Barrett Esophagus) Having Undergone Antireflux Surgery or Not, Compared to the Risk in a Corresponding Background Population, Presented as SIR With 95% CI

No Antireflux Surgery								
Follow-Up, y	Total (n)	Person-Years	Laryngeal or Pharyngeal Squamous Cell Carcinoma		Laryngeal Squamous Cell Carcinoma		Pharyngeal Squamous Cell Carcinoma	
			Cases (n)	SIR (95% CI)	SIR (95% CI)	Cases (n)	Cases (n)	SIR (95% CI)
> 1–34	242,619	1,996,651	321	1.05 (0.94–1.17)	148	1.00 (0.85–1.18)	173	1.09 (0.94–1.27)
> 1–5	242,619	809,131	120	1.09 (0.90–1.30)	55	1.01 (0.76–1.31)	65	1.17 (0.90–1.49)
> 5–10	166,098	640,844	94	0.98 (0.79–1.19)	48	1.04 (0.76–1.37)	46	0.92 (0.67–1.23)
> 10	93,957	546,676	107	1.08 (0.89–1.31)	45	0.97 (0.71–1.30)	62	1.18 (0.91–1.52)
Antireflux Surgery								
Follow-Up, y	Total (n)	Person-Years	Laryngeal or Pharyngeal Squamous Cell Carcinoma		Laryngeal Squamous Cell Carcinoma		Pharyngeal Squamous Cell Carcinoma	
			Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)	Cases (n)	SIR (95% CI)
> 1–34	34,766	425,331	28	0.60 (0.40–0.87)	12	0.54 (0.28–0.94)	16	0.67 (0.38–1.09)
> 1–5	34,766	132,336	5	0.47 (0.15–1.09)	2	0.38 (0.04–1.36)	3	0.56 (0.11–1.62)
> 5–10	31,086	138,999	11	0.78 (0.39–1.39)	7	1.03 (0.41–2.12)	4	0.54 (0.14–1.39)
> 10	23,400	153,996	12	0.56 (0.29–0.98)	3	0.29 (0.06–0.85)	9	0.81 (0.37–1.53)

DISCUSSION

This study indicates that patients with GERD who have undergone antireflux surgery have a decreased risk of laryngeal and pharyngeal squamous cell carcinoma compared to the corresponding background population and to nonoperated patients with GERD (vastly using antireflux medication).

Methodological strengths of this study include the population-based design, encompassing virtually all patients with documented GERD and antireflux surgery in any of the 5 Nordic countries, making it the largest antireflux surgery cohort available to date. This design counteracted selection bias, allowed complete and long follow-up (up to 34 years), provided a large sample size, and facilitated generalizability. The assessment of patients with objectively determined GERD (reflux esophagitis or Barrett esophagus) reduced misclassification of GERD, and the similar results indicate validity of the findings. Another advantage was

the comparison with both the background population and nonoperated patients with GERD, showing similar results.

Possible limitations include residual or unmeasured confounding. The main risk factors for laryngeal and pharyngeal squamous cell carcinoma, tobacco smoking, and alcohol abuse¹⁴ are relevant confounders in this study because tobacco smoking is a risk factor also for GERD, although modest, and alcohol consumption can precipitate reflux episodes among patients with GERD.⁴ Obesity is a risk factor for GERD,⁴ but there is no established association between obesity and laryngeal and pharyngeal squamous cell carcinoma.²³ These 3 potential confounders were adjusted for in the Cox regression analyses. However, because no information on tobacco smoking and alcohol consumption from the participants was available in the registries, we were left to use diagnoses closely associated with these exposures as proxies. Although this strategy could be questioned, it has been

TABLE 4. Risk of Laryngeal and Pharyngeal Squamous Cell Carcinoma Among Patients With Gastroesophageal Reflux Disease Having Undergone Antireflux Surgery or Not, Presented as HR With 95% CI

Follow-Up, y	No Antireflux Surgery		Antireflux Surgery		
	Cases (n)	HR (95% CI)	Cases (n)	Crude HR (95% CI)	Adjusted* HR (95% CI)
Laryngeal or pharyngeal squamous cell carcinoma					
> 1–34	699	1.00 (Reference)	39	0.48 (0.34–0.66)	0.55 (0.38–0.80)
> 1–5	305	1.00 (Reference)	8	0.36 (0.18–0.73)	0.36 (0.16–0.82)
> 5–10	212	1.00 (Reference)	17	0.68 (0.41–1.11)	0.92 (0.53–1.60)
> 10	182	1.00 (Reference)	14	0.40 (0.23–0.69)	0.47 (0.26–0.85)
Laryngeal squamous cell carcinoma					
> 1–34	311	1.00 (Reference)	20	0.55 (0.35–0.87)	0.53 (0.31–0.90)
> 1–5	133	1.00 (Reference)	5	0.52 (0.21–1.27)	0.43 (0.15–1.25)
> 5–10	99	1.00 (Reference)	11	0.94 (0.51–1.76)	1.17 (0.57–2.41)
> 10	79	1.00 (Reference)	4	0.26 (0.10–0.71)	0.23 (0.08–0.69)
Pharyngeal squamous cell carcinoma					
> 1–34	388	1.00 (Reference)	19	0.42 (0.26–0.66)	0.58 (0.35–0.96)
> 1–5	172	1.00 (Reference)	3	0.24 (0.08–0.75)	0.30 (0.08–1.09)
> 5–10	113	1.00 (Reference)	6	0.45 (0.20–1.02)	0.69 (0.29–1.64)
> 10	103	1.00 (Reference)	10	0.52 (0.27–0.99)	0.72 (0.36–1.46)

*Adjusted for sex, age (continuous), calendar period, country, chronic obstructive pulmonary disease, obesity (including diabetes mellitus type 2), and history of excessive alcohol consumption.

TABLE 5. Risk of Laryngeal and Pharyngeal Squamous Cell Carcinoma Among Patients With Severe Gastroesophageal Reflux Disease (With Reflux Esophagitis or Barrett Esophagus) Having Undergone Antireflux Surgery or Not, Presented as HR With 95% CI

Follow-Up, y	No Antireflux Surgery		Antireflux Surgery		
	Cases (n)	HR (95% CI)	Cases (n)	Crude HR (95% CI)	Adjusted* HR (95% CI)
Laryngeal or pharyngeal squamous cell carcinoma					
> 1–34	321	1.00 (Reference)	28	0.48 (0.34–0.66)	0.57 (0.37–0.89)
> 1–5	120	1.00 (Reference)	5	0.36 (0.18–0.73)	0.29 (0.10–0.83)
> 5–10	94	1.00 (Reference)	11	0.68 (0.41–1.11)	1.08 (0.55–2.13)
> 10	107	1.00 (Reference)	12	0.40 (0.23–0.69)	0.54 (0.28–1.04)
Laryngeal squamous cell carcinoma					
> 1–34	148	1.00 (Reference)	12	0.55 (0.35–0.87)	0.42 (0.21–0.83)
> 1–5	55	1.00 (Reference)	2	0.52 (0.21–1.27)	0.23 (0.05–1.14)
> 5–10	48	1.00 (Reference)	7	0.94 (0.51–1.76)	1.16 (0.48–2.84)
> 10	45	1.00 (Reference)	3	0.26 (0.10–0.71)	0.19 (0.05–0.69)
Pharyngeal squamous cell carcinoma					
> 1–34	173	1.00 (Reference)	16	0.42 (0.26–0.66)	0.75 (0.43–1.30)
> 1–5	65	1.00 (Reference)	3	0.24 (0.08–0.75)	0.36 (0.09–1.41)
> 5–10	46	1.00 (Reference)	4	0.45 (0.20–1.02)	0.98 (0.35–2.77)
> 10	62	1.00 (Reference)	9	0.52 (0.27–0.99)	0.95 (0.46–1.98)

*Adjusted for sex, age (continuous), calendar period, country, chronic obstructive pulmonary disease, obesity (including diabetes mellitus type 2) and history of excessive alcohol consumption.

shown to be valid.^{24,25} Because obesity is heavily under-reported in the registries, we added type 2 diabetes as a proxy due to its strong association with obesity, a method of adjusting for obesity that has been shown to be useful.^{25,26} Another limitation is the potential influence of recurrence of GERD following antireflux surgery, which occurred in 17.7% of Swedish participants in the present cohort.²⁷ However, this bias should only attenuate the risk estimates and counteract time-dependent risk reductions over time after antireflux surgery, and not explain the associations. Subclassification of types of antireflux surgery was not possible due to lack of specific surgical codes, but the efficacy on GERD is similar across the main antireflux surgery techniques.^{28–31} In the Nordic countries, the prevalence of GERD is higher than in most developing countries, but comparable to most other Western countries,^{1,2} which suggests that the results could be generalized to Western populations. Finally, despite the large cohort size, the low incidence of laryngeal and pharyngeal squamous cell carcinoma (n=39, 0.08%) reduced the statistical precision, particularly of some subgroup analyses.

The finding of a decreased risk of laryngeal and pharyngeal squamous cell carcinoma following antireflux surgery suggests that such surgery can counteract these tumors in GERD patients as hypothesized. An alternative explanation is biased selection of more healthy and fit individuals for antireflux surgery who may be less prone to be heavy users of tobacco and alcohol, and thus have a lower risk of these tumors than the corresponding background population and nonoperated GERD patients. However, the distribution of diagnoses associated with tobacco smoking and alcohol abuse was similar in GERD patients with and without antireflux surgery, and the results were adjusted for these exposures. Furthermore, the sensitivity analysis of colon cancer, which is unrelated to GERD and antireflux surgery, did not indicate presence of such bias. Additionally, we found no decreased risk of smoking-associated lung cancer over time after antireflux surgery (in manuscript, data not published), and a recent meta-analysis found that GERD increases the risk of laryngeal cancer regardless of history of tobacco smoking and alcohol consumption.¹¹ Finally, the further decreased point estimates of laryngeal squamous cell carcinoma over time after antireflux surgery are unlikely to be explained by selection bias. No such trend over time was found for pharyngeal squamous cell carcinoma, indicating a

need for a more cautious interpretation of the association between antireflux surgery and the risk of this tumor.

Some studies have documented the efficacy of antireflux surgery in the treatment of laryngopharyngeal reflux,³² but no other study has, to our knowledge, investigated whether antireflux surgery prevents laryngeal or pharyngeal cancer. In the aerodigestive tract, conditions causing chronic and repeated injury and inflammation have been linked to cancer development, for example, chronic rhinosinusitis and nasopharyngeal carcinoma,³³ esophagitis and esophageal adenocarcinoma,^{34,35} inflammatory bowel disease and colorectal cancer,³⁶ and chronic hepatitis and hepatocellular carcinoma.³⁷ GERD has been associated with an increased risk of laryngeal and pharyngeal cancer in several studies.^{5,7–12} The lack of association between GERD and laryngeal and pharyngeal squamous cell carcinoma in the present study was unexpected, but could be due to a higher rate of tobacco smoking cessation and alcohol intake reduction in patients who received a GERD diagnosis during in-hospitalization or in specialist outpatient care compared to individuals in the general background population. It is also possible that the frequent use of antireflux medication, which was used in the vast majority of these patients, had some cancer-preventive effects. Moreover, SIRs were dilated toward null association because the high prevalence of persons with GERD diagnosed by general practitioners or other non-specialist outpatient clinics in the general background population were not possible to exclude from the analyses. In vitro, the hypopharyngeal mucosa has shown inflammatory changes similar to the premalignant condition Barrett esophagus following exposure to duodenogastric reflux.¹¹ The laryngeal and pharyngeal epithelium seems to be particularly vulnerable to inflammatory damage following contact with duodenogastric contents, including hydrochloric acid, pepsin, and bile-salts.^{6,38–41} Thus, it is plausible that the hindering of acidic and nonacidic duodenogastric reflux by antireflux surgery would prevent laryngeal and pharyngeal cancer development, whereas antireflux medication, predominant in nonoperated GERD-patients, would be less preventive since it does not hinder duodenogastric reflux, but only reduces its acidic component and the activation of pepsin.⁴² The findings of the present study support this hypothesis.

Considering the low incidence of laryngeal and pharyngeal squamous cell carcinoma,⁴³ the findings of the present study do not suggest recommendation of antireflux surgery for only cancer

protective reasons even if future research confirms the results. However, the findings may become useful for a limited group of high-risk patients for these tumors, and they contribute to the understanding of tumor etiology and suggest higher efficacy of antireflux surgery than antireflux medication in the treatment of patients with more proximal reflux and regurgitation.

In conclusion, this population-based cohort study with long and complete follow-up encompassing all 5 Nordic countries suggests that antireflux surgery in GERD patients decreases the risk of laryngeal squamous cell carcinoma and possibly also pharyngeal squamous cell carcinoma. The results from this first study on the topic need confirmation in future research.

REFERENCES

- Ness-Jensen E, Lindam A, Lagergren J, et al. Changes in prevalence, incidence and spontaneous loss of gastro-oesophageal reflux symptoms: a prospective population-based cohort study, the HUNT study. *Gut*. 2012;61:1390–1397.
- El-Serag HB, Sweet S, Winchester CC, et al. Update on the epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut*. 2014;63:871–880.
- Vakil N, van Zanten SV, Kahrilas P, et al. The Montreal definition and classification of gastroesophageal reflux disease: a global evidence-based consensus. *Am J Gastroenterol*. 2006;101:1900–1920. quiz 1943.
- Ness-Jensen E, Lagergren J. Tobacco smoking, alcohol consumption and gastro-oesophageal reflux disease. *Best Pract Res Clin Gastroenterol*. 2017;31:501–508.
- Lagergren J, Lindam A. Increased risk of laryngeal and pharyngeal cancer after gastrectomy for ulcer disease in a population-based cohort study. *Br J Cancer*. 2012;106:1342–1345.
- Asaoka D, Nagahara A, Matsumoto K, et al. Current perspectives on reflux laryngitis. *Clin J Gastroenterol*. 2014;7:471–475.
- El-Serag HB, Hepworth EJ, Lee P, et al. Gastroesophageal reflux disease is a risk factor for laryngeal and pharyngeal cancer. *Am J Gastroenterol*. 2001;96:2013–2018.
- Zhang D, Zhou J, Chen B, et al. Gastroesophageal reflux and carcinoma of larynx or pharynx: a meta-analysis. *Acta Otolaryngol (Stockh)*. 2014;134:982–989.
- Kuo C-L, Chen Y-T, Shiao A-S, et al. Acid reflux and head and neck cancer risk: a nationwide registry over 13 years. *Auris Nasus Larynx*. 2015;42:401–405.
- Langevin SM, Michaud DS, Marsit CJ, et al. Gastric reflux is an independent risk factor for laryngopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev*. 2013;22:1061–1068.
- Parsel SM, Wu EL, Riley CA, et al. Gastroesophageal and laryngopharyngeal reflux associated with laryngeal malignancy: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. 2019;17:1253–1264. e5.
- Kim SY, Park B, Lim H, et al. Increased risk of larynx cancer in patients with gastroesophageal reflux disease from a national sample cohort. *Clin Otolaryngol*. 2019;44:534–540.
- Lagergren J, Smyth E, Cunningham D, et al. Oesophageal cancer. *Lancet Lond Engl*. 2017;390:2383–2396.
- Rettig EM, D'Souza G. Epidemiology of head and neck cancer. *Surg Oncol Clin N Am*. 2015;24:379–396.
- Maret-Ouda J, Wahlin K, Artama M, et al. Cohort profile: the Nordic Antireflux Surgery Cohort (NordASCo). *BMJ Open*. 2017;7:e016505.
- Maret-Ouda J, Tao W, Wahlin K, et al. Nordic registry-based cohort studies: possibilities and pitfalls when combining Nordic registry data. *Scand J Public Health*. 2017;45:14–19.
- Schmidt M, Schmidt SAJ, Sandegaard JL, et al. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol*. 2015;7:449–490.
- Ludvigsson JF, Andersson E, Ekbom A, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health*. 2011;11:450.
- Sund R. Quality of the Finnish Hospital Discharge Register: a systematic review. *Scand J Public Health*. 2012;40:505–515.
- Pukkala E, Engholm G, Hojsgaard Schmidt LK, et al. Nordic Cancer Registries—an overview of their procedures and data comparability. *Acta Oncol Stockh Swed*. 2018;57:440–455.
- Helweg-Larsen K. The Danish register of causes of death. *Scand J Public Health*. 2011;39:26–29.
- Andersson TM-L, Weiderpass E, Engholm G, et al. Avoidable cancer cases in the Nordic countries—the impact of overweight and obesity. *Eur J Cancer*. 2017;79:106–118.
- Maasland DHE, van den Brandt PA, Kremer B, et al. Body mass index and risk of subtypes of head-neck cancer: the Netherlands Cohort Study. *Sci Rep*. 2015;5:17744.
- Tramacere I, La Vecchia C, Negri E. Tobacco smoking and esophageal and gastric cardia adenocarcinoma: a meta-analysis. *Epidemiol Camb Mass*. 2011;22:344–349.
- Lindblad M, Rodriguez LAG, Lagergren J. Body mass, tobacco and alcohol and risk of esophageal, gastric cardia, and gastric non-cardia adenocarcinoma among men and women in a nested case-control study. *Cancer Causes Control*. 2005;16:285–294.
- Hampel H, Abraham NS, El-Serag HB. Meta-analysis: obesity and the risk for gastroesophageal reflux disease and its complications. *Ann Intern Med*. 2005;143:199–211.
- Maret-Ouda J, Wahlin K, El-Serag HB, et al. Association between laparoscopic antireflux surgery and recurrence of gastroesophageal reflux. *JAAMA*. 2017;318:939–946.
- Roks DJ, Koetje JH, Oor JE, et al. Randomized clinical trial of 270° posterior versus 180° anterior partial laparoscopic fundoplication for gastroesophageal reflux disease. *Br J Surg*. 2017;104:843–851.
- Broeders JA, Roks DJ, Ahmed Ali U, et al. Laparoscopic anterior 180-degree versus nissen fundoplication for gastroesophageal reflux disease: systematic review and meta-analysis of randomized clinical trials. *Ann Surg*. 2013;257:850–859.
- Broeders JA, Roks DJ, Ahmed Ali U, et al. Laparoscopic anterior versus posterior fundoplication for gastroesophageal reflux disease: systematic review and meta-analysis of randomized clinical trials. *Ann Surg*. 2011;254:39–47.
- Broeders F, Mauritz FA, Ahmed Ali U, et al. Systematic review and meta-analysis of laparoscopic Nissen (posterior total) versus Toupet (posterior partial) fundoplication for gastro-oesophageal reflux disease. *Br J Surg*. 2010;97:1318–1330.
- Lechien JR, Dapri G, Dequanter D, et al. Surgical treatment for laryngopharyngeal reflux disease: a systematic review. *JAMA Otolaryngol Head Neck Surg*. 2019;145:655–666.
- Riley CA, Marino MJ, Hawkey N, et al. Sinonasal tract inflammation as a precursor to nasopharyngeal carcinoma: a systematic review and meta-analysis. *Otolaryngol Head Neck Surg*. 2016;154:810–816.
- Lagergren J, Bergström R, Lindgren A, et al. Symptomatic gastroesophageal reflux as a risk factor for esophageal adenocarcinoma. *N Engl J Med*. 1999;340:825–831.
- Rubenstein JH, Taylor JB. Meta-analysis: the association of oesophageal adenocarcinoma with symptoms of gastro-oesophageal reflux. *Aliment Pharmacol Ther*. 2010;32:1222–1227.
- Annese V, Beaugerie L, Egan L, et al. European evidence-based consensus: inflammatory bowel disease and malignancies. *J Crohns Colitis*. 2015;9:945–965.
- Luedde T, Schwabe RF. NF-κB in the liver—linking injury, fibrosis and hepatocellular carcinoma. *Nat Rev Gastroenterol Hepatol*. 2011;8:108–118.
- Sasaki CT, Doukas SG, Costa J, et al. Biliary reflux as a causal factor in hypopharyngeal carcinoma: new clinical evidence and implications. *Cancer*. 2019;125:3554–3565.
- Johnston N, Ondrey F, Rosen R, et al. Airway reflux. *Ann N Y Acad Sci*. 2016;1381:5–13.
- Johnston N, Bulmer D, Gill GA, et al. Cell biology of laryngeal epithelial defenses in health and disease: further studies. *Ann Otol Rhinol Laryngol*. 2003;112:481–491.
- Johnston N, Wells CW, Samuels TL, et al. Pepsin in nonacidic refluxate can damage hypopharyngeal epithelial cells. *Ann Otol Rhinol Laryngol*. 2009;118:677–685.
- Vela MF, Camacho-Lobato L, Srinivasan R, et al. Simultaneous intraesophageal impedance and pH measurement of acid and nonacid gastroesophageal reflux: effect of omeprazole. *Gastroenterology*. 2001;120:1599–1606.
- Ferlay J, Colombet M, Soerjomataram I, et al. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *Int J Cancer*. 2019;144:1941–1953.