



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Different strategies for using topical corticosteroids in people with eczema (Protocol)

Chalmers JR, Axon E, Harvey J, Santer M, Ridd MJ, Lawton S, Langan S, Roberts A, Ahmed A, Muller I, Long CM, Panda S, Chernyshov P, Carter B, Williams HC, Thomas KS

Chalmers JR, Axon E, Harvey J, Santer M, Ridd MJ, Lawton S, Langan S, Roberts A, Ahmed A, Muller I, Long CM, Panda S, Chernyshov P, Carter B, Williams HC, Thomas KS.

Different strategies for using topical corticosteroids in people with eczema.

*Cochrane Database of Systematic Reviews* 2019, Issue 6. Art. No.: CD013356.

DOI: [10.1002/14651858.CD013356](https://doi.org/10.1002/14651858.CD013356).

[www.cochranelibrary.com](http://www.cochranelibrary.com)

Different strategies for using topical corticosteroids in people with eczema (Protocol)

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

---

**TABLE OF CONTENTS**

|                                |    |
|--------------------------------|----|
| HEADER .....                   | 1  |
| ABSTRACT .....                 | 1  |
| BACKGROUND .....               | 2  |
| OBJECTIVES .....               | 4  |
| METHODS .....                  | 4  |
| ACKNOWLEDGEMENTS .....         | 8  |
| REFERENCES .....               | 9  |
| APPENDICES .....               | 12 |
| CONTRIBUTIONS OF AUTHORS ..... | 13 |
| DECLARATIONS OF INTEREST ..... | 13 |
| SOURCES OF SUPPORT .....       | 14 |

---

[Intervention Protocol]

# Different strategies for using topical corticosteroids in people with eczema

Joanne R Chalmers<sup>1</sup>, Emma Axon<sup>1</sup>, Jane Harvey<sup>1</sup>, Miriam Santer<sup>2</sup>, Matthew J Ridd<sup>3</sup>, Sandra Lawton<sup>4</sup>, Sinéad Langan<sup>5</sup>, Amanda Roberts<sup>6</sup>, Amina Ahmed<sup>7</sup>, Ingrid Muller<sup>2</sup>, Chiau Ming Long<sup>8</sup>, Saumya Panda<sup>9</sup>, Pavel Chernyshov<sup>10</sup>, Ben Carter<sup>11</sup>, Hywel C Williams<sup>1</sup>, Kim S Thomas<sup>1</sup>

<sup>1</sup>Centre of Evidence Based Dermatology, University of Nottingham, Nottingham, UK. <sup>2</sup>Department of Primary Care and Population Sciences, University of Southampton, Southampton, UK. <sup>3</sup>University of Bristol, Bristol, UK. <sup>4</sup>Department of Dermatology, Rotherham NHS Foundation Trust, Rotherham, UK. <sup>5</sup>London School of Hygiene and Tropical Medicine, London, UK. <sup>6</sup>Nottingham Support Group for Carers of Children with Eczema, Nottingham, UK. <sup>7</sup>c/o Cochrane Skin Group, University of Nottingham, Nottingham, UK. <sup>8</sup>Department of Pharmacy, School of Medicine, University of Tasmania, Hobart, Australia. <sup>9</sup>Department of Dermatology, KPC Medical College and Hospital, Kolkata, India. <sup>10</sup>Department of Dermatology and Venereology, National Medical University, Kiev, Ukraine. <sup>11</sup>Biostatistics and Health Informatics, King's College London; Institute of Psychiatry, Psychology & Neuroscience, London, UK

**Contact address:** Joanne R Chalmers, Centre of Evidence Based Dermatology, University of Nottingham, Room A103, King's Meadow Campus, Lenton Lane, Nottingham, NG7 2NR, UK. [Joanne.Chalmers@nottingham.ac.uk](mailto:Joanne.Chalmers@nottingham.ac.uk).

**Editorial group:** Cochrane Skin Group.

**Publication status and date:** New, published in Issue 6, 2019.

**Citation:** Chalmers JR, Axon E, Harvey J, Santer M, Ridd MJ, Lawton S, Langan S, Roberts A, Ahmed A, Muller I, Long CM, Panda S, Chernyshov P, Carter B, Williams HC, Thomas KS. Different strategies for using topical corticosteroids in people with eczema. *Cochrane Database of Systematic Reviews* 2019, Issue 6. Art. No.: CD013356. DOI: [10.1002/14651858.CD013356](https://doi.org/10.1002/14651858.CD013356).

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

This is a protocol for a Cochrane Review (Intervention). The objectives are as follows:

To establish the effectiveness and safety of different ways of using topical corticosteroids in people with eczema.

## BACKGROUND

### Description of the condition

Eczema (also called 'atopic dermatitis' or 'atopic eczema') is a chronic inflammatory skin condition, characterised by dry itchy skin with eczematous lesions, which typically fluctuates between periods of remission and flares. Eczema often occurs concurrently with atopic diseases including asthma, allergic rhinitis/hay fever and food allergy. These diseases share a common pathogenesis, and frequently present together in the same individual and family. 'Atopy' refers to the genetic tendency to produce immunoglobulin E (IgE) antibodies in response to small amounts of common environmental proteins such as pollen, house dust mite, and food allergens (Stone 2002; Thomsen 2015). Around 30% of people with eczema develop asthma and 35% develop allergic rhinitis (Luoma 1983). However, it is known that atopy does not concurrently occur in all patients with atopic eczema. In view of this, there have been recent proposals to use the term 'eczema' to define patients both with and without atopy. In agreement with the 'Revised nomenclature for allergy for global use' (Johansson 2004), and for consistency with other Cochrane Reviews evaluating eczema therapies (van Zuuren 2017), we will use the term 'eczema' throughout the review.

Eczema is a common condition throughout the world affecting approximately one in five children, and up to 5% of adults (Barbarot 2018; Odhiambo 2009). The incidence of eczema is highest in the first year of life and can often resolve during childhood (Ban 2018; Kim 2016). However, a recent review has shown that persistence into adolescence and early adulthood may be more common than previously thought, particularly for those with persistent and severe disease or late onset disease (Abuabara 2018).

Eczema can have a significant impact on quality of life (Eckert 2017); and there is a high burden associated with eczema when compared to other skin diseases (Hay 2014). Both the individual and their family can be affected by the disease through factors including disturbed sleep due to itching and scratching, time off work or school for frequent visits to healthcare professionals, restrictions to daily activities, and the need to apply daily time-consuming treatments (Drucker 2016; Eckert 2018).

### Clinical features

Eczema may be acute (short and severe) with weeping vesicles on red, swollen skin, or it may be chronic (long-term) with inflammation, lichenification (thickening of the skin caused by repeated rubbing or scratching), excoriation (abrasion because of rubbing or scratching), hyperpigmentation, and exaggerated surface markings (Weidinger 2016). The typical distribution and type of lesions vary during different stages of life and between different ethnicities. In infants, the extremities and face are usually affected. By around two years of age, lesions mainly appear on the limbs, particularly in the creases of the elbows and knees, as well as the neck, wrists, and ankles. In adulthood, the lesions can become more widespread than those seen in childhood (Bieber 2008a).

The severity of eczema can vary enormously, ranging from dry skin with the occasional itchy inflamed patch, to involvement of the whole body with secondary infections. The course of eczema may also vary from a relapsing-remitting one affecting a few areas recurrently to a continuous one with prolonged periods of inflamed

skin covering most of the body (Berke 2012). Itching can induce a vicious cycle of scratching, leading to skin damage, which in turn causes itchiness—often referred to as the 'itch-scratch cycle' (Pavlis 2017).

### Treatment of eczema

There is currently no cure for eczema, so the treatment goal is control of the disease using the wide range of treatments available including emollients (NICE 2007; SIGN 2011). First-line therapy is the daily application of emollients combined with anti-inflammatory therapy. The most commonly used anti-inflammatory therapy is topical corticosteroids, but topical calcineurin inhibitors are also used. These can be combined with bandages and phototherapy for those who do not respond sufficiently to topical treatment alone. Severe eczema may require systemic treatments such as oral ciclosporin, methotrexate or azathioprine. New biologic agents such as dupilumab are now available for cases of eczema that do not respond to other systemic treatments (Snast 2018). Although topical corticosteroids have been the mainstay of eczema treatment for over 60 years, there are still many unanswered questions about how best to use them (Batchelor 2013).

### Description of the intervention

Topical corticosteroids were first introduced in the 1950s when topical hydrocortisone was found to improve various dermatoses (Sulzberger 1952). Since then, a huge number of topical corticosteroids of increased potency have been developed, and are available in various formulations such as creams and ointments. Mometasone furoate is one of the newer generation of products developed with the intention of producing a safer potent topical corticosteroid (Prakash 1998). Topical corticosteroids are all classified by their potency from mild through to very potent, although the classification of potency varies around the world (British National Formulary 2018; World Health Organization (WHO) 1997). The choice of potency to be used is based on age, body site to be treated and severity of eczema. Low- to moderate-potency topical corticosteroids are usually sufficient for mild eczema and are also used on sensitive skin such as the face and flexural areas. Potent or very potent topical corticosteroids are usually used in severe, thick eczematous plaques over thicker skin sites, such as limbs and palmoplantar surfaces. The advice is to use topical corticosteroids, of appropriate potency, once a day until the eczema is controlled, then 'as required' (NICE 2007).

Local side effects of topical corticosteroids include the possibility of skin atrophy (skin thinning), striae (stretch marks) and purpura (discolouration), whilst systemic side effects include hypothalamic pituitary axis (HPA) suppression and growth suppression (Callen 2007). Skin thinning and effects on growth and development have been reported to be the main concerns amongst people using topical corticosteroids (Li 2017). Side effects of topical corticosteroids are thought to be rare in usual practice and are more likely to occur if topical corticosteroids have been used inappropriately, such as continuous use or if potent corticosteroids are applied to areas which have high permeability (e.g. eyelids) (Callen 2007; Nankervis 2016). This inappropriate use could lead to systemic side effects such as hypothalamic pituitary axis suppression or hyperglycaemia (Gilbertson 1998). But despite their relative safety, concerns and confusion about the use of topical corticosteroids amongst people with eczema and the healthcare

professionals who treat them are widespread. Negative beliefs about the use of topical corticosteroids are thought to contribute to poor treatment adherence (Aubert-Wastiaux 2011; Li 2017; Teasdale 2017).

Topical corticosteroids have traditionally been used reactively (in response to a worsening of the eczema) to control inflammation under the skin. They work by reducing skin inflammation by acting on a number of inflammatory pathways. They bind to glucocorticoid intracellular receptors which then results in a number of anti-inflammatory actions. These include: inhibition of phospholipase A2 activity resulting in reduced production of lipid mediators; inhibition of cyclooxygenase induction causing decreased prostaglandin production; inhibition of nitric oxide synthase production; inhibition of cytokine causing suppression of cell-mediated inflammation; inhibition of mast cell activity resulting in decreased levels of mast cell inflammatory mediators; and vasoconstriction (local blood flow reduction) (Ahluwalia 1998).

A number of different ways (or 'strategies') of using topical corticosteroids for treating eczema have been proposed. Proactive use of topical corticosteroids for two days per week between flares is thought to help to prevent eczema flares and therefore reduce the need for more intense periods of topical corticosteroid use to treat flares which may be associated with an increase in adverse events (Schmitt 2011). Applying topical corticosteroids to wet skin after bathing or use of wet wraps may increase penetration through the skin and increase delivery of the cream or ointment into the upper layers, thus increasing efficacy of the topical corticosteroid (Gonzalez-Lopez 2017; Kohn 2016). Topical calcineurin inhibitors (pimecrolimus or tacrolimus) can be used instead of topical corticosteroids and a strategy of alternating between these two treatments may be as effective as using topical corticosteroids alone, but may reduce the adverse events associated with topical corticosteroids, such as skin thinning (Broeders 2016).

Some strategies aim to reduce adverse events whilst increasing or maintaining effectiveness of the topical corticosteroid. Applying topical corticosteroids once daily may be as effective as two or more times a day but may reduce the likelihood of adverse events occurring (Green 2004). Another strategy is to use different potency topical corticosteroids, possibly combined with different duration of use, such as a more potent topical corticosteroid for a shorter period compared to milder potency topical corticosteroids for a longer duration. This reduces the length of time the topical corticosteroid would be used although more potent topical corticosteroids may be associated with increased adverse events (Thomas 2002).

Since topical corticosteroids are used with emollients, other proposed strategies concern the combined use of these two treatments, such as the order in which the treatments are applied and the optimum time lapse between application of each treatment. Current guidance in the UK from the National Health Service (NHS) is to apply emollients first then wait for 30 minutes before applying the topical corticosteroid for maximal benefit (NHS 2019). Additionally, different preparations of topical corticosteroids (e.g. ointments, creams) have been developed to increase the efficacy; and different concentrations (e.g. hydrocortisone 0.5% versus 2.5%). More recently, 'second generation' once-daily topical corticosteroids (mometasone furoate and fluticasone propionate) have been proposed as a safer and effective alternative to the older topical corticosteroid preparations (Bieber 2008b).

## Why it is important to do this review

It is well established that some patients, parents and clinicians have considerable concerns about using topical corticosteroids for treating eczema (Charman 2000; Li 2017; Teasdale 2017). As a result, topical corticosteroids are often under-used in Western countries, resulting in poorly controlled disease (Lundin 2018). Conversely, in other areas of the world, such as India, potent topical corticosteroid use is often unregulated and patients are able to obtain these steroids over the counter. Subsequent inappropriate use of potent topical corticosteroids can lead to an increase in adverse events (Coondoo 2014).

This situation is exacerbated by the lack of clarity as to how the different ways of using topical corticosteroids — such as once-a-day or twice-a-day application, increasing topical corticosteroid potency in response to a flare, or twice-a-week use to proactively prevent flares — affect both effectiveness and safety profile. The British National Formulary (BNF) provides little reassurance, describing adrenal suppression as rare but providing no quantification of other side effects (British National Formulary 2018). Concerns and uncertainties around topical corticosteroids were highlighted in the James Lind Alliance Priority Setting Partnership for eczema, in which the following two topics relating to topical corticosteroid safety were identified by patients and healthcare professionals as priority areas for research (Batchelor 2013).

- “What is the best and safest way of using topical corticosteroids?”
- “What is the long-term safety of topical corticosteroids?”

This comprehensive systematic review is needed to summarise the available evidence on the effectiveness and safety of different ways of using topical corticosteroids to support patients and clinicians in making informed treatment choices. However, since most eczema trials have a relatively short follow-up, this review will primarily address the first of these two questions.

The strategies included in this review will refer to different methods of using topical corticosteroids to improve effectiveness or safety, or both, and hence achieve the best outcomes for the patients. A strategy may aim to improve the long-term control of eczema, for example, in the case of proactive topical corticosteroid treatment. This strategy involves weekly application of topical corticosteroid, for two consecutive days, to previously affected or new sites of eczema, to reduce the risk of flares (Schmitt 2011). Alternatively, a strategy such as reducing the frequency of application may be designed to improve the safety of the drug whilst maintaining effectiveness (Green 2004; Williams 2007).

This review forms part of a body of work funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research (grant no: RP-PG-0216-20007) to develop an online behavioural intervention to support self-care of eczema in children, adolescents and young adults (Eczema Care Online, ECO), and the findings will contribute to development of the intervention by providing data on the best and safest ways to use topical corticosteroids.

This review will be complemented by another ongoing Cochrane Review which will incorporate a network meta-analysis: "Topical treatments for eczema: a network meta-analysis". The ongoing

Cochrane Review will compare topical corticosteroids to other topical treatments, such as topical calcineurin inhibitors.

## OBJECTIVES

To establish the effectiveness and safety of different ways of using topical corticosteroids in people with eczema.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

All randomised controlled trials (RCTs) where randomisation is at any level (including cluster and within-participants trials).

#### Types of participants

Participants of any age and gender with a diagnosis of eczema (also known as atopic dermatitis or atopic eczema) of any severity by a healthcare professional, or using established diagnostic criteria for eczema (e.g. the standardised diagnostic criteria of Hanifin and Rajka (Hanifin 1980) or the UK Working Party diagnostic criteria for atopic dermatitis (Williams 1994)). We will also accept modified versions of standard diagnostic criteria.

Studies which include participants with other types of eczema such as contact dermatitis, varicose eczema, and seborrhoeic eczema, or other inflammatory or 'steroid-responsive' skin conditions such as psoriasis, will only be included if the trial also includes people with eczema and the data are reported separately.

#### Types of interventions

The intervention will be any topical corticosteroid of any preparation and potency in a trial where one strategy of using the topical corticosteroid is compared to a different strategy. This will include, but not necessarily be limited to, the following strategies.

#### Different strategies for topical corticosteroid use

- Proactive versus reactive topical corticosteroid treatment (e.g. twice per week versus 'as required').
- Frequency of application of topical corticosteroid (e.g. once daily versus twice daily).
- Duration of use of topical corticosteroids for induction of remission (e.g. short term versus long term).
- Different time of day for applying topical corticosteroids, and order of application or timing between applications of topical corticosteroids and emollients (e.g. evening versus morning, immediately following a bath/shower versus no guidance).

#### Different preparations or potencies of topical corticosteroids

- Different potencies of topical corticosteroids (e.g. mild versus moderate; stepping up or stepping down potency of topical corticosteroids).
- Different preparations but the same topical corticosteroid (e.g. ointment versus cream), including proprietary topical corticosteroid versus the generic equivalent (e.g. Elocon versus mometasone furoate 0.1%).
- Different concentrations of topical corticosteroids (e.g. hydrocortisone 0.5% versus hydrocortisone 2.5%).

### Topical corticosteroids alongside other interventions

- Alternating between topical corticosteroids and calcineurin inhibitors (e.g. topical corticosteroids alone versus alternating between topical corticosteroids and topical calcineurin inhibitors).
- Use of topical corticosteroids under occlusion (e.g. with wet wraps versus no wet wraps).

### Other strategies

- Second generation topical corticosteroids (mometasone furoate and fluticasone propionate) against first generation topical corticosteroids of the same potency (e.g. mometasone furoate 0.1% versus betamethasone valerate 0.1%).
- A combination of any of the strategies above (e.g. short burst of potent topical corticosteroid versus longer duration of mild topical corticosteroids).
- Any other strategies for using topical corticosteroids not listed above.

Since the focus of this review is to compare different strategies of using topical corticosteroids, the following comparisons will be excluded.

- Topical corticosteroid compared with either no treatment, vehicle or placebo (unless it is specifically assessing an alternative regimen such as proactive therapy).
- Topical corticosteroid compared with another topical corticosteroid of the same potency and preparation but no differences in how they are used.
- Topical corticosteroid compared with different topical treatments such as calcineurin inhibitors or emollients.
- Topical corticosteroid compared with systemic treatments.
- Topical corticosteroid treatment in conjunction with an eczema treatment used for the most severe cases of eczema as defined by NICE (i.e. phototherapy and systemic therapy) (NICE 2018). This is because it will be difficult to detect any differences in efficacy or safety between the topical corticosteroids strategies when such treatments are also used.

### Types of outcome measures

We will assess both effectiveness and safety to reflect the overall aim of this review.

The effectiveness outcomes of interest for this review are focused on the two domains for which the international Harmonizing Outcome Measures for Eczema (HOME) initiative has recommended core outcome measurement instruments i.e. clinician-reported signs and patient-reported symptoms of eczema (HOME).

There is currently no agreed standardised timing for effectiveness outcome assessments for eczema trials. Therefore, to assess treatment effects in a consistent way, we will focus on short-term effectiveness outcomes reported between one and four weeks (taking the earliest available time point within that range), medium-term effectiveness outcomes between 12 and 16 weeks (taking the closest time point to 12 weeks), and long-term effectiveness as the longest time point greater than 16 weeks.

We will also report outcomes at baseline, end of treatment, and end of follow-up regardless of timing. We will attempt to pool data at similar time points where possible.

Because many different instruments are used to assess effectiveness of treatments for eczema (Schmitt 2007), we will use a hierarchical approach in which we will initially extract data from one instrument per outcome based on the priority order described below. However, we will also make a note of other instruments reported, and data from other lower priority instruments may also be extracted to maximize our ability to summarise data in pooled analyses. Where possible, we will also compare the effect sizes calculated in this review against the minimal clinically important differences (MCID) from the literature.

Throughout this review, we will use the term 'effectiveness' to describe both 'efficacy' and 'effectiveness'. In many trials it is likely to be unclear whether the trial is primarily assessing efficacy or effectiveness and we would prefer to avoid making inappropriate judgements.

Safety outcomes of interest reflect the side effects of topical corticosteroids. We define 'relevant' adverse events as those previously identified as being of particular concern to patients (Li 2017), the side effects listed in the Summary of Product Characteristics for topical corticosteroids used to treat eczema, and original data submissions from the Eczema Priority Setting Partnership, outlining patients' and clinicians' concerns about the safety of topical corticosteroids (Batchelor 2013). We will report data on individual relevant adverse events and their relatedness to the study drug where available. Where appropriate, the total number of relevant adverse events per treatment group will be reported to allow pooling of data. We will present safety data for the treatment and follow-up periods separately where possible. Where outcomes are assessed during post-treatment follow-up, we will only include data where participants are retained in their randomised groups. We will not use long-term safety data from crossover or within-participant trials due to the high likelihood of contamination.

For safety data, we will report adverse events at the end of treatment and the end of follow-up (where specified).

### Primary outcomes

Two primary outcomes are included — safety and effectiveness (clinician-reported signs of eczema) — to reflect the overall aim of this review.

- **Changes in clinician-reported signs of eczema (effectiveness).** We will extract data based on the following priority order of instruments.
  - Eczema Area and Severity Index (EASI) – this is the HOME recommended core outcome measurement instrument for clinician-reported signs of eczema (Hanifin 2001;Schmitt 2014).
  - Objective SCORing Atopic Dermatitis (ObjSCORAD) – measures similar aspects of the disease to EASI (Kunz 1997;Oranje 2007).
  - SCORing Atopic Dermatitis (SCORAD) – most commonly used instrument in eczema trials, comprising objective SCORAD plus itch and sleep loss (Kunz 1997;Oranje 2007)

- Six Area, Six Sign Atopic Dermatitis (SASSAD) severity score (Berth-Jones 1996).
- Three Item Severity score (TIS) (Oranje 2007;Willemssen 2009;Wolkerstorfer 1999).
- Investigator Global Assessment (IGA) – no validated instrument and little consistency between trials but commonly included (Futamura 2016).
- Any other instruments.
- **Number of relevant local adverse events (safety).** This will include skin thinning, striae, telangiectasia, aging/wrinkling, changes in skin colour, sensitisation, skin bleaching, worsening or induction of acne, skin infections, folliculitis, perioral/periocular dermatitis, and application site reactions such as burning sensation/stinging. In our analyses, we will primarily focus on the number of participants with at least one adverse event.

### Secondary outcomes

- **Patient-reported symptoms of eczema (effectiveness).** Data will be extracted based on the following priority order of instruments.
  - Patient-Oriented Eczema Measure (POEM) – recommended core instrument by HOME for the patient-reported symptoms of eczema (Charman 2004;Spuls 2017).
  - Patient-Oriented SCORAD (PO-SCORAD) (Vourc'h-Jourdain 2009).
  - Sleep and itch scales, as measured by Visual Analogue Scales (VAS) or Numerical Rating Scales (NRS).
  - Self-Administered EASI (SA-EASI) (Housman 2002).
  - Patient Global Assessment (PGA) – no validated instrument and little consistency between trials.
  - Any other instruments.
- **Number of relevant systemic adverse events (safety).** This will include bone problems, impact on growth and development, effects on endocrine system, eye problems, and cancer. In our analyses, we will primarily focus on the number of participants with at least one adverse event.

### Search methods for identification of studies

We aim to identify all relevant randomised controlled trials (RCTs) regardless of language or publication status (published, unpublished, in press, or in progress).

### Electronic searches

The Cochrane Skin Information Specialist will search the following databases for relevant trials with no restriction by date.

- the Cochrane Skin Specialised Register;
- the Cochrane Central Register of Controlled Trials (CENTRAL) in the Cochrane Library;
- MEDLINE via Ovid (from 1946 onwards);
- Embase via Ovid (from 1974 onwards);
- the GREAT database (Global Resource for Eczema Trials (Centre of Evidence Based Dermatology); [www.greatdatabase.org.uk](http://www.greatdatabase.org.uk)).

The Information Specialist has devised a draft search strategy for RCTs for MEDLINE (Ovid), which is displayed in Appendix 1. We will

use this as the basis for search strategies for the other databases listed.

### **Trial registers**

We (EA, JH) will search the following trial registers using the search terms 'eczema' and 'atopic dermatitis'.

- the ISRCTN register ([www.isrctn.com](http://www.isrctn.com));
- ClinicalTrials.gov ([www.clinicaltrials.gov](http://www.clinicaltrials.gov));
- the Australian New Zealand Clinical Trials Registry ([www.anzctr.org.au](http://www.anzctr.org.au));
- the World Health Organization International Clinical Trials Registry Platform (ICTRP) ([apps.who.int/trialsearch/](http://apps.who.int/trialsearch/)); and
- the EU Clinical Trials Register ([www.clinicaltrialsregister.eu](http://www.clinicaltrialsregister.eu)).

### **Searching other resources**

#### **Searching reference lists**

We will check the bibliographies of included studies and any relevant systematic reviews identified for further references to relevant trials.

#### **Correspondence with trialists**

We will contact trialists for clarification and further data if trial reports are unclear.

#### **Adverse effects**

We will not perform a separate search for adverse effects of the target interventions. We will consider adverse effects described in included studies only.

### **Data collection and analysis**

We will use [Covidence](#) systematic review software to screen and manage the references, and to record the data extracted from the included studies. The software will also automatically create a study flow diagram (PRISMA) for us to include in the review ([Liberati 2009](#)).

#### **Selection of studies**

Two review authors (EA, JH) will independently screen the titles and abstracts of each record identified in the searches. If a study meets our inclusion criteria, we will analyse the full text to confirm its inclusion. Any disagreement will be resolved by a third reviewer (JRC, KST or HCW). We will record reasons for exclusions in the 'Characteristics of excluded studies' table. We will present the process of trial selection in a PRISMA flow diagram ([Liberati 2009](#)).

#### **Data extraction and management**

Two review authors (EA, JH) will independently extract data from each included study using a data extraction form. We will pilot and modify the form, as necessary. The following data will be extracted.

- Study population (e.g. inclusion criteria, setting/country, severity of eczema, age, gender, ethnicity)
- Study methods (e.g. study design, blinding methods, funding source)
- Interventions and comparators (e.g. treatment name, potency, frequency of use)

- Primary and secondary outcomes (including time points – both during treatment and during follow-up)

We will enter study characteristics into a 'Characteristics of included studies' table; and we will analyse or narratively describe extracted outcome data. We will resolve any disagreements during the data extraction phase through discussion with a third author (JRC, KST, or HCW).

We will include multiple reports of a trial but will choose a primary reference and list the others as secondary references.

We will also extract study characteristics for ongoing trials to produce a 'Characteristics of ongoing studies' table. We will do the same for 'Studies awaiting classification' where there is not enough information available for us to include or exclude the trial in the review.

### **Assessment of risk of bias in included studies**

Two authors (EA, JH) will independently assess the risk of bias of each included study using the Cochrane 'Risk of bias' tool ([Higgins 2011](#)). The following domains will be assessed.

- Selection bias (random sequence generation and allocation concealment)
- Performance bias (blinding of participants and study personnel)
- Detection bias (blinding of outcome assessment)
- Attrition bias (completeness of data, missing data and losses to follow-up, intention to treat principle)
- Reporting bias (selective reporting of outcomes, assessed via comparing with the study's protocol or clinical trial register entry)
- Other bias (including design-specific risks of bias, baseline imbalance, contamination, fraud, selective reporting of subgroups)

Each domain will be assessed as low, unclear or high risk of bias. Disagreements will be resolved via discussion with a third author (JRC, KST or HCW). We will present a 'Risk of bias' graph and 'Risk of bias' summary figure in the review.

### **Measures of treatment effect**

We will report dichotomous data as risk ratios (RR) with associated 95% confidence intervals (CI). Continuous data will be reported as mean differences (MD) with associated 95% CIs where the same scale is used to measure an outcome. Where appropriate, we may consider using a standardised mean difference (SMD) and associated 95% CI when different instruments are used to measure effectiveness outcomes (i.e. clinician-reported signs and patient-reported symptoms).

For adverse event data, we will focus on analysing the number of participants with at least one adverse event, and hence we will report RR. However, in some studies the number of events per arm may only be reported and it will not be clear if multiple adverse events occurred within the same individual. For these studies, we will contact the authors to request data in the format of 'number of participants with at least one adverse event'. If we are unable to obtain these data, we will analyse these data as count data, following guidance from the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)), and produce rate ratios

instead of RR. Where studies report time-to-flare data based on a measure of effectiveness (e.g. EASI), we shall extract hazard ratios (HR) from the study reports.

### Unit of analysis issues

The unit of analysis will primarily be the individual participant. If only count data is available for adverse events, we will analyse the number of events over time and calculate rate ratios. We will only meta-analyse cluster RCTs with parallel RCTs if the data reported in the trial publication has been correctly analysed, taking into account the number of clusters. If appropriate, we may try to estimate the intracluster correlation coefficient to enable us to include the trial in the meta-analysis (Higgins 2011).

For within-participant trials, we will extract paired data from the study report and analyse separately to parallel RCTs. If paired data are not available, we will attempt to estimate the paired data using appropriate methods and guidance from the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011; Hirji 2011). If this is not possible, we will only use data from the first part of the study in cross-over trials, or narratively describe the results.

For within-participant trials where body parts receive different interventions, we will not extract data on outcomes which affect the whole body (e.g. systemic adverse events), as it will not be possible to determine which treatment caused the event.

For studies which include multiple intervention groups, we will either analyse each intervention group versus a comparator in a separate analysis, or combine groups to create a single pair-wise comparison, if clinically appropriate (Higgins 2011).

### Dealing with missing data

Where possible we will conduct an intention-to-treat analysis, including all randomised participants where data is provided. If data is missing, we will contact study authors and produce a table in the review detailing such contact (e.g. dates, information requested, whether they replied). We will conduct sensitivity analyses, removing studies at high risk of attrition bias and will also consider attrition bias when undertaking our quality assessments (Schünemann 2013). For continuous data, we will attempt to calculate any missing statistics (e.g. standard deviations) using the methods described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

### Assessment of heterogeneity

When pooling studies in a meta-analysis, we will consider any methodological and clinical differences between studies, and only include studies in the same meta-analysis where it is considered appropriate. We will assess heterogeneity through forest plot inspection and the  $I^2$  statistic using the thresholds defined in the *Cochrane Handbook for Systematic Reviews of Interventions*: 0% to 40% might not be important; 30% to 60% may represent moderate heterogeneity; 50% to 90% may represent substantial heterogeneity; and 75% to 100% represents considerable heterogeneity (Higgins 2011). If we identify substantial or considerable statistical heterogeneity, we will attempt to determine reasons by examining the study characteristics and performing subgroup analyses where appropriate.

### Assessment of reporting biases

If we include 10 or more studies in a meta-analysis, we will produce a funnel plot to explore publication bias (Sterne 2011). We will describe narratively suspected reporting biases and their potential effects on the overall results and conclusions.

### Data synthesis

We will narratively synthesise outcome data and will conduct meta-analysis (if appropriate) in Review Manager 5, using the random-effects model (Review Manager 2014). We will use the generic inverse variance model to analyse count data (rate ratios) and may use it to compare studies that have presented a group difference, or have adjusted analysis only. We will present effect estimates, with 95% confidence intervals and associated  $I^2$  statistic and P value, for all pooled synthesis. For dichotomous outcomes, if statistical evidence of an effect is reported we will calculate an associated number needed to treat for an additional beneficial outcome (with associated 95% confidence intervals).

### Subgroup analysis and investigation of heterogeneity

We plan the following subgroup analyses.

- Children versus adults
- Anatomical site e.g. topical corticosteroid applied to sensitive sites (face/genitals) versus other body sites
- Baseline severity of eczema (mild disease versus moderate and severe, as specified in the trial report)

If there is substantial statistical heterogeneity (via forest plot inspection and using the  $I^2$  statistic), additional clinical and methodological differences will be investigated. Clinical differences may include filaggrin (FLG) mutation status, age sub-groups of children (0 to 4, 5 to 11, and 11+ years), chronic versus acute disease, and body surface area affected.

### Sensitivity analysis

If possible, we will perform a sensitivity analysis removing studies at high risk of bias from the meta-analysis. We will also perform sensitivity analyses where assumptions have been made or data imputed.

### 'Summary of findings' tables and GRADE assessments

We will create 'Summary of findings' tables for our main comparisons. We have selected the following four most relevant and important comparisons, from both clinician and patient perspectives, to be included in the 'Summary of findings' tables.

1. Proactive versus reactive topical corticosteroid treatment (e.g. twice per week versus 'as required')
2. Frequency of application of topical corticosteroid (e.g. once daily versus twice daily)
3. Duration of use of topical corticosteroids for induction of remission (e.g. short term versus long term)
4. Different potencies of topical corticosteroids (e.g. mild versus moderate; stepping up or stepping down topical corticosteroids)

We will include both primary outcomes and secondary outcomes in each 'Summary of findings' table. We will use the GRADE approach to assess the certainty of evidence for each primary and secondary outcome for our main comparisons. GRADE includes

the assessment of five factors: study limitations (risk of bias); inconsistency of results; indirectness of evidence; imprecision; and publication bias (Schünemann 2013). Each outcome can be downgraded by one or two levels for each domain, and we will class the overall certainty as high, moderate, low or very low. We will use GRADEpro to create our 'Summary of findings' tables and undertake our GRADE assessments (GRADEpro GDT).

## ACKNOWLEDGEMENTS

We would like to thank Douglas Grindlay for his input into this protocol.

We would also like to thank Liz Doney, the information specialist at Cochrane Skin, for her assistance in drafting the search strategy.

The Cochrane Skin editorial base wishes to thank Ching-Chi Chi, Cochrane Dermatology Editor; Toby Lasserson, who provided methodological comment; the clinical referee, Yik Weng Yew; and Jason Elliot-Smith, who copy-edited the protocol.

## REFERENCES

### Additional references

#### Abuabara 2018

Abuabara K, Yu AM, Okhovat JP, Allen IE, Langan SM. The prevalence of atopic dermatitis beyond childhood: A systematic review and meta-analysis of longitudinal studies. *Allergy* 2018;**73**(3):696-704.

#### Ahluwalia 1998

Ahluwalia A. Topical glucocorticoids and the skin-mechanisms of action: an update. *Mediators of Inflammation* 1998;**7**(3):183-93.

#### Aubert-Wastiaux 2011

Aubert-Wastiaux H, Moret L, Le Rhun A, Fontenoy AM, Nguyen JM, Leux C, et al. Topical corticosteroid phobia in atopic dermatitis: a study of its nature, origins and frequency. *British Journal of Dermatology* 2011;**165**(4):808-14.

#### Ban 2018

Ban L, Langan SM, Abuabara K, Thomas KS, Abdul Sultan A, Sach T, et al. Incidence and sociodemographic characteristics of eczema diagnosis in children: a cohort study. *Journal of Allergy and Clinical Immunology* 2018;**141**(5):1927-9. [DOI: [10.1016/j.jaci.2017.12.997](https://doi.org/10.1016/j.jaci.2017.12.997)]

#### Barbarot 2018

Barbarot S, Auziere S, Gadkari A, Girolomoni G, Puig L, Simpson EL, et al. Epidemiology of atopic dermatitis in adults: Results from an international survey. *Allergy* 2018;**73**(6):1284-93. [DOI: [10.1111/all.13401](https://doi.org/10.1111/all.13401)]

#### Batchelor 2013

Batchelor JM, Ridd MJ, Clarke T, Ahmed A, Cox M, Crowe S, et al. The Eczema Priority Setting Partnership: a collaboration between patients, carers, clinicians and researchers to identify and prioritize important research questions for the treatment of eczema. *British Journal of Dermatology* 2013;**168**:577-82.

#### Berke 2012

Berke R, Singh A, Guralnick M. Atopic dermatitis: an overview. *American Family Physician* 2012;**86**(1):35-42.

#### Berth-Jones 1996

Berth-Jones J. Six area, six sign atopic dermatitis (SASSAD) severity score: a simple system for monitoring disease activity in atopic dermatitis. *British Journal of Dermatology* 1996;**135** Suppl 48:25-30.

#### Bieber 2008a

Bieber T. Atopic dermatitis. *New England Journal of Medicine* 2008;**358**(14):1483-94.

#### Bieber 2008b

Bieber T. Atopic and contact dermatitis. In: Rich R, Fleisher T, Shearer W, Schroeder H, Frew A, Weyand C editor(s). *Clinical Immunology: Principles and Practice*. 3rd Edition. St Louis, MO: Mosby, 2008:667-80.

#### British National Formulary 2018

Joint Formulary Committee. British National Formulary (online). [www.bnf.nice.org.uk](http://www.bnf.nice.org.uk) (accessed 28 June 2018).

#### Broeders 2016

Broeders JA, Ahmed Ali U, Fischer G. Systematic review and meta-analysis of randomized clinical trials (RCTs) comparing topical calcineurin inhibitors with topical corticosteroids for atopic dermatitis: A 15-year experience. *Journal of the American Academy of Dermatology* 2016;**75**(2):410-9.e3.

#### Callen 2007

Callen J, Chamlin S, Eichenfield LF, Ellis C, Girardi M, Goldfarb M, et al. A systematic review of the safety of topical therapies for atopic dermatitis. *British Journal of Dermatology* 2007;**156**(2):203-21.

#### Charman 2000

Charman CR, Morris AD, Williams HC. Topical corticosteroid phobia in patients with atopic eczema. *British Journal of Dermatology* 2000;**142**(5):931-6.

#### Charman 2004

Charman CR, Venn AJ, Williams HC. The patient-oriented eczema measure: development and initial validation of a new tool for measuring atopic eczema severity from the patients' perspective. *Archives of Dermatology* 2004;**140**(12):1513-9.

#### Coondoo 2014

Coondoo A. Topical corticosteroid misuse: the Indian scenario. *Indian Journal of Dermatology* 2014;**59**(5):451-5.

#### Covidence [Computer program]

Veritas Health Innovation. Covidence. Melbourne, Australia: Veritas Health Innovation, 2017.

#### Drucker 2016

Drucker AM, Wang AR, Qureshi AA. Research gaps in quality of life and economic burden of atopic dermatitis: the National Eczema Association burden of disease audit. *JAMA Dermatology* 2016;**152**(8):873-4.

#### Eckert 2017

Eckert L, Gupta S, Amand C, Gadkari A, Mahajan P, Gelfand JM. Impact of atopic dermatitis on health-related quality of life and productivity in adults in the United States: An analysis using the National Health and Wellness Survey. *Journal of the American Academy of Dermatology* 2017;**77**(2):274-9. e3.

#### Eckert 2018

Eckert L, Gupta S, Amand C, Gadkari A, Mahajan P, Gelfand JM. The burden of atopic dermatitis in US adults: Health care resource utilization data from the 2013 National Health and Wellness Survey. *Journal of the American Academy of Dermatology* 2018;**78**(1):54-61. e1.

#### Futamura 2016

Futamura M, Leshem YA, Thomas KS, Nankervis H, Williams HC, Simpson EL. A systematic review of Investigator Global

Assessment (IGA) in atopic dermatitis (AD) trials: Many options, no standards. *Journal of the American Academy of Dermatology* 2016;**74**(2):288-94.

#### **Gilbertson 1998**

Gilbertson EO, Spellman MC, Piacquadio DJ, Mulford MI. Super potent topical corticosteroid use associated with adrenal suppression: clinical considerations. *Journal of the American Academy of Dermatology* 1998;**38**(2 Pt 2):318-21.

#### **Gonzalez-Lopez 2017**

Gonzalez-Lopez G, Ceballos-Rodriguez RM, Gonzalez-Lopez JJ, Feito Rodriguez M, Herranz-Pinto P. Efficacy and safety of wet wrap therapy for patients with atopic dermatitis: a systematic review and meta-analysis. *British Journal of Dermatology* 2017;**177**(3):688-95.

#### **GRADEpro GDT [Computer program]**

McMaster University (developed by Evidence Prime). GRADEpro GDT. Hamilton (ON): McMaster University (developed by Evidence Prime), 2015.

#### **Green 2004**

Green C, Colquitt JL, Kirby J, Davidson P, Payne E. Clinical and cost-effectiveness of once-daily versus more frequent use of same potency topical corticosteroids for atopic eczema: a systematic review and economic evaluation. *Health Technology Assessment* 2004;**8**(47):iii,iv, 1-120.

#### **Hanifin 1980**

Hanifin JM, Rajka G. Diagnostic features of atopic dermatitis. *Acta Dermato-venereologica* 1980;**92**:44-7.

#### **Hanifin 2001**

Hanifin JM, Thurston M, Omoto M, Cherill R, Tofte SJ, Graeber M. The eczema area and severity index (EASI): assessment of reliability in atopic dermatitis. EASI Evaluator Group. *Experimental Dermatology* 2001;**10**(1):11-8.

#### **Hay 2014**

Hay RJ, Williams NE, Williams HC, Bolliger IW, Dellavalle RP, Margolis DJ, et al. The global burden of skin disease in 2010: an analysis of the prevalence and impact of skin conditions. *Journal of Investigative Dermatology* 2014;**134**(6):1527-34.

#### **Higgins 2011**

Higgins JP, Green S, editor(s). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

#### **Hirji 2011**

Hirji KF, Fagerland MW. Calculating unreported confidence intervals for paired data. *BMC Medical Research Methodology* 2011;**11**:66.

#### **HOME**

Harmonising Outcome Measures for Eczema (HOME). [www.homeforeczema.org/](http://www.homeforeczema.org/) (accessed prior to 03 May 2018).

#### **Housman 2002**

Housman TS, Patel MJ, Camacho F, Feldman SR, Fleischer AB Jr, Balkrishnan R. Use of the Self-Administered Eczema Area and Severity Index by parent caregivers: results of a validation study. *British Journal of Dermatology* 2002;**147**(6):1192-8.

#### **Johansson 2004**

Johansson SG, Bieber T, Dahl R, Friedmann PS, Lanier BQ, Lockey RF, et al. Revised nomenclature for allergy for global use: Report of the Nomenclature Review Committee of the World Allergy Organization October 2003. *Journal of Allergy and Clinical Immunology* 2004;**113**(5):832-6. [PUBMED: 15131563]

#### **Kim 2016**

Kim JP, Chao LX, Simpson EL, Silverberg JI. Persistence of atopic dermatitis (AD): A systematic review and meta-analysis. *Journal of American Academy of Dermatology* 2016;**75**(4):681-687.e11.

#### **Kohn 2016**

Kohn LL, Kang Y, Antaya RJ. A randomized, controlled trial comparing topical steroid application to wet versus dry skin in children with atopic dermatitis (AD). *Journal of the American Academy of Dermatology* 2016;**75**(2):306-11.

#### **Kunz 1997**

Kunz B, Oranje AP, Labreze L, Stalder JF, Ring J, Taieb A. Clinical validation and guidelines for the SCORAD index: consensus report of the European Task Force on Atopic Dermatitis. *Dermatology* 1997;**195**(1):10-9.

#### **Li 2017**

Li AW, Yin ES, Antaya RJ. Topical corticosteroid phobia in atopic dermatitis: A systematic review. *JAMA Dermatology* 2017;**153**(10):1036-42.

#### **Liberati 2009**

Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Medicine* 2009;**6**(7):e1000100.

#### **Lundin 2018**

Lundin S, Wahlgren CF, Bergstrom A, Johansson EK, Dahlen E, Andersson N, et al. Use of emollients and topical glucocorticoids among adolescents with eczema - data from the population-based birth cohort BAMSE. *British Journal of Dermatology* 2018;**179**(3):709-16.

#### **Luoma 1983**

Luoma R, Koivikko A, Viander M. Development of asthma, allergic rhinitis and atopic dermatitis by the age of five years. A prospective study of 543 newborns. *Allergy* 1983;**38**(5):339-46. [PUBMED: 6614406]

#### **Nankervis 2016**

Nankervis H, Thomas KS, Delamere FM, Barbarot S, Rogers NK, Williams HC. Scoping systematic review of treatments for eczema. Southampton (UK): NIHR Journals Library. Programme Grants for Applied Research, 2016.

**NHS 2019**

National Health Service (NHS). Topical corticosteroids. [www.nhs.uk/conditions/topical-steroids/](http://www.nhs.uk/conditions/topical-steroids/) (accessed 05 February 2019).

**NICE 2007**

NICE. Clinical guideline [CG57]. Atopic eczema in under 12s: diagnosis and management 2007. [www.nice.org.uk/Guidance/CG57](http://www.nice.org.uk/Guidance/CG57) (accessed prior to 03 May 2018).

**NICE 2018**

NICE. Eczema overview. Treating atopic eczema in children aged 12 and under. [pathways.nice.org.uk/pathways/eczema](http://pathways.nice.org.uk/pathways/eczema) (accessed 26 July 2018).

**Odhiambo 2009**

Odhiambo JA, Williams HC, Clayton TO, Robertson CF, Asher MI, ISAAC Phase Three Study Group. Global variations in prevalence of eczema symptoms in children from ISAAC Phase Three. *Journal of Allergy and Clinical Immunology* 2009;**124**(6):1251-1258.e23.

**Oranje 2007**

Oranje AP, Glazenburg EJ, Wolkerstorfer A, de Waard-van der Spek FB. Practical issues on interpretation of scoring atopic dermatitis: the SCORAD index, objective SCORAD and the three-item severity score. *British Journal of Dermatology* 2007;**157**(4):645-8.

**Pavlis 2017**

Pavlis J, Yosipovitch G. Management of itch in atopic dermatitis. *American Journal of Clinical Dermatology* 2017 [Epub ahead of print]. [DOI: [10.1007/s40257-017-0335-4](https://doi.org/10.1007/s40257-017-0335-4).]

**Prakash 1998**

Prakash A, Benfield P. Topical mometasone. A review of its pharmacological properties and therapeutic use in the treatment of dermatological disorders. *Drugs* 1998;**55**(1):145-63.

**Review Manager 2014 [Computer program]**

Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

**Schmitt 2007**

Schmitt J, Langan S, Williams HC, European Dermato-Epidemiology Network. What are the best outcome measurements for atopic eczema? A systematic review. *Journal of Allergy and Clinical Immunology* 2007;**120**(6):1389-98.

**Schmitt 2011**

Schmitt J, Von Kobyletzki L, Svensson A, Apfelbacher C. Efficacy and tolerability of proactive treatment with topical corticosteroids and calcineurin inhibitors for atopic eczema: Systematic review and meta-analysis of randomized controlled trials. *British Journal of Dermatology* 2011;**164**(2):415-28.

**Schmitt 2014**

Schmitt J, Spuls PI, Thomas KS, Simpson E, Furue M, Deckert S, et al. The Harmonising Outcome Measures for Eczema (HOME)

statement to assess clinical signs of atopic eczema in trials. *Journal of Allergy and Clinical Immunology* 2014;**134**(4):800-7.

**Schünemann 2013**

Schünemann H, Brozek J, Oxman A, editors. GRADE handbook for grading quality of evidence and strength of recommendation. Updated October 2013. [www.guidelinedevelopment.org/handbook](http://www.guidelinedevelopment.org/handbook). The GRADE Working Group. Available from: (accessed prior to 03 May 2018).

**SIGN 2011**

SIGN (Scottish Intercollegiate Guidelines Network). Management of atopic eczema in primary care 2011. [www.sign.ac.uk](http://www.sign.ac.uk) (accessed prior to 03 May 2018).

**Snast 2018**

Snast I, Reiter O, Hodak E, Friedland R, Mimouni D, Leshem YA. Are Biologics Efficacious in Atopic Dermatitis? A Systematic Review and Meta-Analysis. *American Journal of Clinical Dermatology* 2018;**19**(2):145-65.

**Spuls 2017**

Spuls PI, Gerbens LAA, Simpson E, Apfelbacher CJ, Chalmers JR, Thomas KS, et al. Patient-Oriented Eczema Measure (POEM), a core instrument to measure symptoms in clinical trials: a Harmonising Outcome Measures for Eczema (HOME) statement. *British Journal of Dermatology* 2017;**176**(4):979-84.

**Sterne 2011**

Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011;**343**:d4002.

**Stone 2002**

Stone KD. Atopic diseases of childhood. *Current Opinion in Pediatrics* 2002;**14**(5):634-46. [PUBMED: 12352260]

**Sulzberger 1952**

Sulzberger MB, Witten VH. The effect of topically applied compound F in selected dermatoses. *Journal of Investigative Dermatology* 1952;**19**(2):101-2.

**Teasdale 2017**

Teasdale EJ, Muller I, Santer M. Carers' views of topical corticosteroid use in childhood eczema: a qualitative study of online discussion forums. *British Journal of Dermatology* 2017;**176**(6):1500-7.

**Thomas 2002**

Thomas KS, Armstrong S, Avery A, Po AL, O'Neill C, Young S, et al. Randomised controlled trial of short bursts of a potent topical corticosteroid versus prolonged use of a mild preparation for children with mild or moderate atopic eczema. *BMJ* 2002;**324**(7340):768.

**Thomsen 2015**

Thomsen SF. Epidemiology and natural history of atopic diseases. *European Clinical Respiratory Journal* 2015;**2**(1):24642. [PUBMED: 26557262]

**van Zuuren 2017**

van Zuuren EJ, Fedorowicz Z, Christensen R, Lavrijsen A, Arents BWM. Emollients and moisturisers for eczema. *Cochrane Database of Systematic Reviews* 2017, Issue 2. [DOI: [10.1002/14651858.CD012119.pub2](https://doi.org/10.1002/14651858.CD012119.pub2)]

**Vourc'h-Jourdain 2009**

Vourc'h-Jourdain M, Barbarot S, Taieb A, Diepgen T, Ambonati M, Durosier V, et al. Patient-oriented SCORAD: a self-assessment score in atopic dermatitis. A preliminary feasibility study. *Dermatology* 2009;**218**(3):246-51.

**Weidinger 2016**

Weidinger S, Novak N. Atopic dermatitis. *Lancet* 2016;**387**(10023):1109-22.

**Willemsen 2009**

Willemsen MG, van Valburg RW, Dirven-Meijer PC, Oranje AP, van der Wouden JC, Moed H. Determining the severity of atopic dermatitis in children presenting in general practice: an easy and fast method. *Dermatology Research and Practice* 2009;**2009**:357046.

**Williams 1994**

Williams HC, Burney PG, Hay RJ, Archer CB, Shipley MJ, Hunter JJ, et al. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. I. Derivation of a minimum set of discriminators for atopic dermatitis. *British Journal of Dermatology* 1994;**131**(3):383-96.

**Williams 2007**

Williams HC. Established corticosteroid creams should be applied only once daily in patients with atopic eczema. *BMJ* 2007;**334**(7606):1272.

**Wolkerstorfer 1999**

Wolkerstorfer A, de Waard van der Spek FB, Glazenburg EJ, Mulder PG, Oranje AP. Scoring the severity of atopic dermatitis: three item severity score as a rough system for daily practice and as a pre-screening tool for studies. *Acta Dermatovenereologica* 1999;**79**(5):356-9.

**World Health Organization (WHO) 1997**

World Health Organization (WHO). WHO model prescribing information: drugs used in skin diseases. [www.apps.who.int/medicinedocs/en/d/Jh2918e/32.html](http://www.apps.who.int/medicinedocs/en/d/Jh2918e/32.html) (accessed 6 July 2018).

## APPENDICES

### Appendix 1. Draft MEDLINE (Ovid) search strategy

1. exp Eczema/ or eczema\$.ti,ab.
2. exp Dermatitis, Atopic/
3. neurodermatitis.ti,ab. or exp Neurodermatitis/
4. exp Dermatitis/ or dermatitis.ti,ab.
5. besnier\$ prurigo.ti,ab.
6. prurigo diathesique.ti,ab.
7. or/1-6
8. (topical\$ adj3 corticosteroid\$.ti,ab.
9. (topical\$ adj3 steroid\$.ti,ab.
10. (topical\$ adj3 corticoid\$.ti,ab.
11. (topical\$ adj3 glucocorticoid\$.ti,ab.
12. exp Desonide/
13. desonide.mp.
14. alclometasone.mp.
15. amcinonide.mp.
16. exp Beclomethasone/
17. (beclometasone or beclomethasone).mp.
18. exp Betamethasone/
19. Betamethasone.mp.
20. budesonide.mp. or exp Budesonide/
21. clobetasol\$.mp. or exp Clobetasol/
22. clobetasone\$.mp.
23. clocortolone.mp.
24. (exp Cortisone/ or cortisone.ti,ab.) and (exp Administration, Topical/ or exp Ointments/ or Dermatologic Agents/)
25. Deprodone.mp.
26. desoximetasone.mp. or exp Desoximetasone/
27. exp Dexamethasone/ or dexamethasone.mp.
28. dichlorisone.mp.
29. diflorasone.mp.
30. exp Diflucortolone/ or diflucortolone.mp.
31. Difluprednate.mp.
32. fluclorolone.mp.
33. Flucloronide.mp.
34. Fludrocortisone.mp.

35. fludroxycortide.mp.
36. (flumetasone or flumethasone).mp.
37. exp Flumethasone/
38. fluocinolone.mp.
39. fluocinonide.mp. or exp Fluocinonide/
40. fluocortin.mp.
41. exp Fluocortolone/
42. fluocortolone.mp.
43. Fluorometholone.mp.
44. fluprednidene.mp.
45. flurandrenolide.mp.
46. flurandrenolone.mp. or exp Flurandrenolone/
47. fluticasone.mp.
48. halcinonide.mp. or exp Halcinonide/
49. halobetasol.mp.
50. halometasone.mp.
51. exp Hydrocortisone/
52. cortisol.ti.ab. and (exp Administration, Topical/ or exp Ointments/ or Dermatologic Agents/)
53. hydrocortisone\$.mp.
54. (masipredone or Mazipredone).mp.
55. exp Methylprednisolone/
56. methylprednisolone.mp.
57. mometasone.mp.
58. prednicarbat\$.mp.
59. exp Prednisolone/
60. (Prednisolone or prednisone).mp.
61. ulobetasol.mp.
62. triamcinolone.mp. or exp Triamcinolone/
63. exp Adrenal Cortex Hormones/ and (exp Administration, Topical/ or exp Ointments/ or Dermatologic Agents/)
64. exp Glucocorticoids/ and (exp Administration, Topical/ or exp Ointments/ or Dermatologic Agents/)
65. or/8-64
66. randomized controlled trial.pt.
67. controlled clinical trial.pt.
68. randomized.ab.
69. placebo.ab.
70. clinical trials as topic.sh.
71. randomly.ab.
72. trial.ti.
73. 66 or 67 or 68 or 69 or 70 or 71 or 72
74. exp animals/ not humans.sh.
75. 73 not 74
76. 7 and 65 and 75

[Lines 66-75: Cochrane Highly Sensitive Search Strategy for identifying randomized trials in MEDLINE: sensitivity- and precision-maximizing version (2008 revision)]

## CONTRIBUTIONS OF AUTHORS

JRC was the contact person with the editorial base.

JRC, EA, and JH co-ordinated the contributions from the co-authors and wrote the final draft of the protocol.

JRC, EA, JH, and BC worked on the Methods section.

JRC, EA, and JH drafted the clinical sections of the Background and responded to the clinical comments of the referees.

JRC, EA, JH, and BC responded to the methodology and statistics comments of the referees.

All authors contributed to writing the protocol.

AR and AA were the consumer co-authors and checked the protocol for readability and clarity. They also ensured that the outcomes are relevant to consumers.

## DECLARATIONS OF INTEREST

JRC: member of the HOME executive, ECO co-applicant

EA: none known

JH: none known

KST: member of the HOME executive, ECO joint-lead-applicant

**Different strategies for using topical corticosteroids in people with eczema (Protocol)**

Copyright © 2019 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

MS: none known

HCW: Chair of the HOME executive, Developed the POEM scale, ECO co-applicant

BC: none known

SL: ECO co-applicant, Honorarium Thorton Ross – lecture

MR: funded by National Institute for Health Research (NIHR) Post-doctoral Fellowship (PDF-2014-07-013)

SML: funded by Wellcome Senior Clinical fellowship

CML: none known.

SP: Merck, FDC & GSK-Stiefel (who produce products that may be used in atopic eczema) are advertisers in the Indian Journal of Dermatology, Venereology & Leprology of which I am the Editor-in-Chief.

PC: none known.

AR: ECO co-applicant

AA: ECO co-applicant

IM: ECO co-applicant

### **Disclaimer**

This project was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to the Cochrane Skin Group. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS or the Department of Health.

### **SOURCES OF SUPPORT**

#### **Internal sources**

- No sources of support supplied

#### **External sources**

- The National Institute for Health Research (NIHR), UK.

The NIHR, UK, is the largest single funder of the Cochrane Skin Group.

- NIHR, UK.

This review forms part of a body of work funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research (grant no: RP-PG-0216-20007) to develop an online behavioural intervention to support self-care of atopic eczema in children, adolescents and young adults and the findings will contribute to development of the intervention by providing data on the best and safest ways to use topical corticosteroids.